

**From the Cover: Low-dose alcohol actions on  $\alpha$ 4 $\beta$ 3 $\delta$  GABAA receptors are reversed by the behavioral alcohol antagonist Ro15-4513**

M. Wallner, H. J. Hanchar, and R. W. Olsen

*PNAS* 2006;103;8540-8545; originally published online May 12, 2006;  
doi:10.1073/pnas.0600194103

**This information is current as of April 2007.**

<b>Online Information &amp; Services</b>	High-resolution figures, a citation map, links to PubMed and Google Scholar, etc., can be found at: <a href="http://www.pnas.org/cgi/content/full/103/22/8540">www.pnas.org/cgi/content/full/103/22/8540</a>
<b>Related Articles</b>	A related article has been published: <a href="http://www.pnas.org/cgi/content/full/103/22/8307">www.pnas.org/cgi/content/full/103/22/8307</a>
<b>References</b>	This article cites 67 articles, 22 of which you can access for free at: <a href="http://www.pnas.org/cgi/content/full/103/22/8540#BIBL">www.pnas.org/cgi/content/full/103/22/8540#BIBL</a>  This article has been cited by other articles: <a href="http://www.pnas.org/cgi/content/full/103/22/8540#otherarticles">www.pnas.org/cgi/content/full/103/22/8540#otherarticles</a>
<b>E-mail Alerts</b>	Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or <a href="#">click here</a> .
<b>Rights &amp; Permissions</b>	To reproduce this article in part (figures, tables) or in entirety, see: <a href="http://www.pnas.org/misc/rightperm.shtml">www.pnas.org/misc/rightperm.shtml</a>
<b>Reprints</b>	To order reprints, see: <a href="http://www.pnas.org/misc/reprints.shtml">www.pnas.org/misc/reprints.shtml</a>

Notes:

# Low-dose alcohol actions on $\alpha 4\beta 3\delta$ GABA<sub>A</sub> receptors are reversed by the behavioral alcohol antagonist Ro15-4513

M. Wallner\*<sup>†</sup>, H. J. Hanchar\*, and R. W. Olsen<sup>†</sup>

Department of Molecular and Medical Pharmacology, University of California, Los Angeles, CA 90095-1735

Edited by Richard L. Huganir, Johns Hopkins University School of Medicine, Baltimore, MD, and approved April 7, 2006 (received for review January 9, 2006)

Although it is now more than two decades since it was first reported that the imidazobenzodiazepine Ro15-4513 reverses behavioral alcohol effects, the molecular target(s) of Ro15-4513 and the mechanism of alcohol antagonism remain elusive. Here, we show that Ro15-4513 blocks the alcohol enhancement on recombinant "extrasynaptic"  $\alpha 4/6\beta 3\delta$  GABA<sub>A</sub> receptors at doses that do not reduce the GABA-induced Cl<sup>-</sup> current. At low ethanol concentrations ( $\leq 30$  mM), the Ro15-4513 antagonism is complete. However, at higher ethanol concentrations ( $\geq 100$  mM), there is a Ro15-4513-insensitive ethanol enhancement that is abolished in receptors containing a point mutation in the second transmembrane region of the  $\beta 3$  subunit ( $\beta 3N265M$ ). Therefore,  $\alpha 4/6\beta 3\delta$  GABA receptors have two distinct alcohol modulation sites: (i) a low-dose ethanol site present in  $\alpha 4/6\beta 3\delta$  receptors that is antagonized by the behavioral alcohol antagonist Ro15-4513 and (ii) a site activated at high (anesthetic) alcohol doses, defined by mutations in membrane-spanning regions. Receptors composed of  $\alpha 4\beta 3N265M\delta$  subunits that lack the high-dose alcohol site show a saturable ethanol dose-response curve with a half-maximal enhancement at 16 mM, close to the legal blood alcohol driving limit in most U.S. states (17.4 mM). Like in behavioral experiments, the alcohol antagonist effect of Ro15-4513 on recombinant  $\alpha 4\beta 3\delta$  receptors is blocked by flumazenil and  $\beta$ -carboline-ethyl ester ( $\beta$ -CCE). Our findings suggest that ethanol/Ro15-4513-sensitive GABA<sub>A</sub> receptors are important mediators of behavioral alcohol effects.

alcohol intoxication | alcohol receptor | anesthetics

Although alcohol is one of the most widely used and abused drugs, the molecular targets that mediate alcohol effects at concentrations relevant for mild social intoxication are only beginning to be revealed. Neurotransmitter receptors for GABA, NMDA and glycine, and G protein-gated K<sup>+</sup> channels have been identified as potential alcohol targets that are sensitive to intoxicating alcohol concentrations (1–4). GABA<sub>A</sub> receptors (GABA<sub>A</sub>Rs) have long been suspected to be important mediators of alcohol effects (5, 6) because benzodiazepines (BZs) and barbiturates, classic GABA<sub>A</sub>R agonists, share common pharmacological properties with ethanol, such as sedative-hypnotic, anti-anxiety, and motor in-coordinating and anticonvulsant effects and have additive, possibly even synergistic effects, when taken together with ethanol (7). In addition, BZs, barbiturates, and ethanol produce tolerance and cross-tolerance to each other (8), consistent with GABA<sub>A</sub>Rs as targets of action.

We have recently identified subtypes of GABA<sub>A</sub>Rs, those containing the  $\delta$  and the  $\beta 3$  subunit, that are uniquely sensitive to low alcohol concentrations (9). Consistent with the view that  $\delta$  subunit-containing receptors are important mediators of alcohol actions is the finding that GABA<sub>A</sub>R  $\delta$ -subunit knockout mice show multiple defects in behavioral responses to ethanol (10). Receptors containing the  $\delta$  subunit have an exclusively nonsynaptic distribution, are sensitive to low ambient extrasynaptic GABA concentrations, and display slow desensitization,

properties that enable them to mediate a persistent (or tonic) form of inhibition (11). GABA<sub>A</sub>R  $\delta$  subunits seem to almost exclusively associate with the GABA<sub>A</sub>R  $\alpha 4$  and  $\alpha 6$  subunits, two closely related and somewhat specialized  $\alpha$  subunits that confer insensitivity to classical BZ agonists (but not the imidazobenzodiazepines flumazenil and Ro15-4513), because they carry the amino acid arginine ( $\alpha 6R100$ ,  $\alpha 4R100$  instead of histidine present in  $\alpha 1$ , -2, -3, and -5) at a site critical for BZ binding (12). We showed that the  $\alpha 6$ -R100Q "BZ-site" mutation, previously identified in alcohol-nontolerant rats (13), is sufficient for behavioral alcohol hypersensitivity, confers alcohol supersensitivity to tonic currents in rat cerebellar granule cells, and dramatically increases the alcohol sensitivity of recombinantly expressed receptors, if the  $\alpha 6R100Q$  subunit is expressed together with the  $\beta 3$  and the  $\delta$  subunit (14). Therefore, an amino acid residue important for BZ sensitivity is also critical for low-dose alcohol sensitivity of  $\alpha 6$ -containing GABA<sub>A</sub>R.

Two decades ago it was reported that the imidazobenzodiazepine Ro15-4513, a structural analog of the clinically used general BZ antagonist flumazenil, is an alcohol antagonist in mammals (15, 16). The reversal of motor-impairing, sedative, anxiolytic, amnestic, and rewarding alcohol actions by Ro15-4513 at low to moderate alcohol doses were confirmed by various groups in different species (17–24). However, the debate whether Ro15-4513's alcohol antagonism is due to a specific alcohol counteracting action, or due to nonspecific functional antagonism, due to the weak inverse agonist activity of Ro15-4513 on certain GABA<sub>A</sub>R subtypes (12), so far has not been resolved (18, 19, 25–28). Three observations argue against a nonspecific, inverse agonist action of Ro15-4513: (i) other inverse agonists are not alcohol antagonists (17, 18); (ii) GABA<sub>A</sub>R-mediated Cl<sup>-</sup> flux in synaptoneurosomes is enhanced by ethanol (17, 29–31) and this enhancement can be blocked by Ro15-4513 at concentrations that do not inhibit the GABA current (17); and (iii) in experiments where Ro15-4513 leads to a partial reversal of sedative-hypnotic alcohol actions, it did not reduce the hypothermic actions (32). Hypothermic and analgesic alcohol actions are at least partially mediated by G protein-gated potassium channels (3, 4, 33, 34).

Here, we show that the enhancement of  $\delta$  subunit-containing recombinant GABA<sub>A</sub>R at low to moderately intoxicating ethanol doses (3–30 mM) is antagonized by the behavioral alcohol antagonist Ro15-4513, suggesting that these subtypes of

Conflict of interest statement: M.W., R.W.O. and H.J.H. have filed a U.S. Provisional Patent Application, Serial No. 60/693,844.

This paper was submitted directly (Track II) to the PNAS office.

Abbreviations:  $\beta$ -CCE,  $\beta$ -carboline-3-carboxy ethyl ester; GABA<sub>A</sub>R, GABA<sub>A</sub> receptor; DMCM, methyl-6,7-dimethoxy-4-ethyl- $\beta$ -carboline-3-carboxylate; BZ, benzodiazepine.

See Commentary on page 8307.

\*M.W. and H.J.H. contributed equally to this work.

<sup>†</sup>To whom correspondence may be addressed. E-mail: mwallner@mednet.ucla.edu or rolsen@mednet.ucla.edu.


© 2006 by The National Academy of Sciences of the USA

GABA<sub>A</sub>Rs, previously thought to be completely insensitive to BZs, have a high-affinity Ro15-4513-binding site. At anesthetic and potentially lethal alcohol concentrations ( $\geq 100$  mM), there is an additional Ro15-4513-insensitive component of ethanol enhancement. We show that this component disappears in  $\alpha 4\beta 3\delta$  receptors that contain a mutation in the  $\beta 3$  subunit ( $\beta 3N265M$ ). This mutation has been previously shown to essentially abolish (in  $\beta 3N265M$  knockin mice) the *in vivo* anesthetic actions of etomidate and propofol (35), consistent with the identification of this residue in determining the  $\beta$  subunit selectivity for enhancement of recombinant GABA<sub>A</sub>Rs by loreclezole and etomidate (36, 37). Homologous residues in GABA<sub>A</sub>R subunits have been identified as important for high-dose alcohol and volatile anesthetic actions on recombinant GABA<sub>A</sub>Rs (38). As expected for a “true” receptor,  $\alpha 4\beta 3N265M\delta$  GABA<sub>A</sub>Rs that lack the Ro15-4513-insensitive high-dose ethanol enhancement now show a saturable alcohol dose-response curve with a half-maximal effect at 16 mM. As previously shown in behavioral assays, Ro15-4513’s alcohol antagonism on recombinant receptors can be abolished by flumazenil. Most importantly, the fact that low-dose alcohol, as well as the BZ alcohol antagonist Ro15-4513, can exert its effects on common GABA<sub>A</sub>R subtypes identifies ethanol/Ro15-4513-sensitive GABA<sub>A</sub> receptors as important mediators of alcohol intoxication at low to moderate alcohol concentrations.

## Results


**Low-Dose Ethanol Enhancement on  $\alpha 4/6\beta 3\delta$  GABA<sub>A</sub>Rs Is Antagonized by Ro15-4513.** Inspired by reports that Ro15-4513 antagonizes most low-dose ethanol actions in animals (17, 19, 32) and that 100 nM Ro15-4513 blocks the low-dose (30 mM) alcohol enhancement in GABA-induced  $^{36}\text{Cl}^-$  flux in brain homogenate assays (17, 39), we decided to investigate the effect of the BZ alcohol antagonist Ro15-4513 on the alcohol-induced current enhancement in  $\alpha 4/6\beta 3\delta$  receptors expressed in oocytes. Fig. 1 shows that 300 nM Ro15-4513 completely reversed the ethanol enhancement of  $\alpha 4\beta 3\delta$  GABA<sub>A</sub>Rs for ethanol concentrations up to 30 mM. This “anti-alcohol effect” of Ro15-4513 is surprisingly specific because, at concentrations where it abolished the alcohol-induced current increase (up to 300 nM), Ro15-4513 did not lead to a reduction in the “basal” GABA-induced current on these receptors; i.e., it is not an inverse agonist in this assay at concentrations that inhibit the ethanol-augmentation of GABA currents (Fig. 1c). The dose dependence of this effect revealed that the concentration of Ro15-4513 required to inhibit 50% of the 30 mM ethanol enhancement ( $\text{IC}_{50}$ ) was  $\approx 10$  nM (Fig. 1b). The alcohol antagonist action of low-dose Ro15-4513 suggests that, against common knowledge,  $\alpha 4/6\beta 3\delta$  GABA<sub>A</sub>R have a high-affinity Ro15-4513-binding site, with a  $K_d$  of  $\approx 10$  nM.

At higher alcohol concentrations ( $\geq 100$  mM), a fraction of the alcohol-induced enhancement was not blocked by 300 nM Ro15-4513 (Fig. 1a). This high-dose ethanol enhancement was not surmountable by increasing the Ro15-4513 concentrations (data not shown). Therefore,  $\alpha 4\beta 3\delta$  GABA<sub>A</sub>Rs have a distinct Ro15-4513-insensitive component of alcohol enhancement. To demonstrate the dose dependence, we applied increasing concentrations of Ro15-4513 to currents evoked by 300 nM GABA plus 10, 30, or 50 mM ethanol. Ro15-4513 led to a dose-dependent block that was complete for the 10- and 30-mM dose (with 300 nM Ro15-4513) (Fig. 1b and c). Again, at the 50-mM ethanol dose, a small fraction of the alcohol enhancement was not blocked by Ro15-4513 (Fig. 1c). The complete and specific antagonism of low-dose alcohol effects on these receptors by Ro15-4513 suggests the intriguing possibility that Ro15-4513 might work by competitively displacing EtOH from its binding site.



**Fig. 1.** Ro15-4513 antagonizes ethanol effects on recombinant  $\alpha 4\beta 3\delta$  receptor currents. (a and b) To mimic tonic GABA current, 300 nM GABA was perfused onto *Xenopus* oocytes expressing rat  $\alpha 4\beta 3\delta$  receptors that were held at  $-80$  mV, and currents were measured with a two-electrode voltage clamp system. The indicated doses of ethanol and drugs were applied. (a) Ro15-4513 (300 nM) completely antagonized ethanol enhancement up to an ethanol concentration of 30 mM. (b) Cumulative Ro15-4513 dose-response curve shows a dose-dependent inhibition of ethanol effects. GABA-evoked currents were blocked by  $30 \mu\text{M}$  bicuculline. (c) GABA peak currents with and without ethanol and the indicated concentrations of Ro15-4513. Ro15-4513 led to a dose-dependent inhibition of (10, 30, and 50 mM) ethanol enhancement on  $\alpha 4\beta 3\delta$  receptors. At concentrations up to 300 nM, Ro15-4513 did not block the basal current on  $\alpha 4\beta 3\delta$  receptors (evoked by 300 nM GABA). (a and b) Representative recordings of six and five experiments, respectively.

**Antagonizing Ro15-4513’s Alcohol Antagonism by Flumazenil and  $\beta$ -Carboline-Ethyl Ester ( $\beta$ -CCE).** Certain BZ site ligands, like the general BZ antagonist flumazenil (Ro15-1788) and the structurally unrelated BZ-site ligands  $\beta$ -CCE and FG7142, were shown to prevent the alcohol antagonist effects of Ro15-4513 in behavioral assays (17, 39). We reasoned that this result could be due to displacement of Ro15-4513 from its binding site by these compounds, which do not show alcohol antagonism by themselves (17, 39). We therefore tested four selected BZ site ligands for their ability to reverse or mimic Ro15-4513 antagonism of ethanol effects. We applied these compounds to  $\alpha 4\beta 3\delta$  receptors (expressed in oocytes) in the presence of 300 nM ( $\approx \text{EC}_{20}$ ) GABA (to mimic tonic GABA current), 30 mM ethanol (to increase the GABA current), and 100 nM Ro15-4513 (to reverse the enhancement by 30 mM ethanol). Flumazenil (Ro15-1788) and  $\beta$ -CCE at 300 nM reversed the Ro15-4513-induced alcohol antagonism. However, the classical BZ agonist flunitrazepam, as well as DMCM (methyl-6,7-dimethoxy-4-ethyl- $\beta$ -carboline-3-carboxylate) (a  $\beta$ -carboline with pronounced inverse agonist efficacy on  $\gamma 2$  subunit-containing receptors), each at  $1 \mu\text{M}$ , did not reverse the effects of Ro15-4513 (Fig. 2a), indicating that the Ro15-4513/BZ-binding site on  $\delta$  subunit-containing receptors is unique and binds only certain BZs and BZ site ligands with high affinity. None of the four compounds tested blocked ethanol




**Fig. 2.** Ro15-4513 alcohol antagonism is antagonized by flumazenil and  $\beta$ -CCE, but not flunitrazepam and DMCM. (a) Currents were evoked by 300 nM GABA and potentiated by 30 mM ethanol, and this potentiation was reversed by 100 nM Ro15-4513. In constant presence of 300 nM GABA, 30 mM ethanol, and 100 nM Ro15-4513, the BZ site ligands Ro15-1788 (300 nM),  $\beta$ -CCE (300 nM), flunitrazepam, and DMCM (each 1  $\mu$ M) were sequentially applied to test whether they reverse Ro15-4513's ethanol antagonist effects. At the end of the recording, 30  $\mu$ M bicuculline was used to block the GABA-induced current. Shown is a representative recording of a total of three experiments. (b) Chemical structures of the imidazobenzodiazepines Ro15-4513 and Ro15-1788 show that they differ only at one single position in the molecule. The clinically used BZ antagonist flumazenil (Ro15-1788) carries a fluorine at the C7 position of the BZ ring, whereas Ro15-4513 carries the larger azido group.

enhancement on their own (data not shown). These data are consistent with previous findings that Ro15-4513's alcohol antagonism can be antagonized by certain BZ-site ligands in  $^{36}\text{Cl}^-$  flux assays in synaptoneurosomes and provide an *in vitro* correlate to the behavioral data that show that flumazenil and  $\beta$ -CCE can reverse the alcohol antagonist effects of Ro15-4513 (17, 39). A comparison of the structures of Ro15-4513 and flumazenil shows that these two molecules are identical, except for one moiety, which is an azido group in Ro15-4513 and a fluorine in flumazenil (Fig. 2b).

**$\beta$ -CCE Is a Positive GABA Modulator on  $\alpha 4\beta 3\delta$  Receptors.** We consistently observed that  $\beta$ -CCE led to an "overshoot" when we used it to antagonize the alcohol antagonist effects of Ro15-4513 on GABA/ethanol-induced currents (see Fig. 2), suggesting that  $\beta$ -CCE might potentiate alcohol effects on  $\alpha 4/6\beta 3\delta$  GABA<sub>A</sub>R. We therefore tested  $\beta$ -CCE alone and in combination with 3 mM ethanol for their functional effects on  $\alpha 4\beta 3\delta$  receptors. Fig. 3a shows that  $\beta$ -CCE not only enhances alcohol actions, but also increases the activity of  $\alpha 4\beta 3\delta$  GABA<sub>A</sub>Rs in the absence of alcohol. In the same way as alcohol effects on  $\alpha 4\beta 3\delta$  GABA<sub>A</sub>R, the  $\beta$ -CCE-induced enhancement of GABA currents is reversed by Ro15-4513 (Fig. 3b). A likely explanation for these findings is that  $\beta$ -CCE, as well as Ro15-4513, occupies an overlapping and mutually exclusive binding site, whereas  $\beta$ -CCE and ethanol might be able to bind next to each other in a side-by-side binding pocket, both microdomains blocked by Ro15-4513 (see Fig. 2b).

**Loss of Ro15-4513-Insensitive Ethanol Actions in  $\alpha 4\beta 3\text{N}265\text{M}\delta$  GABA<sub>A</sub>R.** The Ro15-4513-insensitive component of ethanol enhancement is observed at high alcohol concentrations (>30 mM), where most recombinant and native GABA<sub>A</sub>R show ethanol enhancement that is likely due to alcohol sites determined by




**Fig. 3.**  $\beta$ -CCE enhances ethanol effects and is an agonist on  $\alpha 4\beta 3\delta$  receptors. (a) The  $\beta$ -carboline  $\beta$ -CCE at the indicated concentrations was applied alone or together with ethanol (always in the presence of 300 nM GABA) to oocytes expressing  $\alpha 4\beta 3\delta$  receptors, and peak GABA/ $\text{Cl}^-$  currents were measured. (b) Dose-dependent reversal of 300 nM  $\beta$ -CCE enhancement of  $\alpha 4\beta 3\delta$  currents by Ro15-4513.

mutations in the second and third transmembrane region of GABA<sub>A</sub>Rs (38). We show here (Fig. 4) that  $\alpha 4\beta 3\text{N}265\text{M}\delta$  receptors, where the  $\beta 3$  WT subunit is replaced with the mutated  $\beta 3\text{N}265\text{M}$  subunit, retain the Ro15-4513-sensitive alcohol enhancement. However, the  $\beta 3\text{N}265\text{M}$  mutation completely abolished the Ro15-4513-insensitive ethanol enhancement observed at 100 and 300 mM ethanol (Fig. 4a), and even at 1 M ethanol (data not shown). GABA<sub>A</sub>R composed of  $\alpha 4\beta 3\text{N}265\text{M}\delta$  and  $\alpha 4\beta 3\delta$  subunits show identical ethanol enhancement at alcohol concentrations up to 30 mM and differ only at the 100- and 300-mM dose (Fig. 4b). As a consequence of this loss of Ro15-4513-insensitive, high-concentration alcohol effects, recombinant  $\alpha 4\beta 3\text{N}265\text{M}\delta$  receptors now have a saturable ethanol response curve with a half-maximal response of 16 mM [at  $\text{EC}_{50}$  (300 nM) GABA, Fig. 2b], a concentration close to the legal blood alcohol limit (17 mM) for driving in most US states.

## Discussion

**Extrasynaptic  $\delta$  Subunit-Containing Receptors as Targets for Ethanol Action.** GABA<sub>A</sub>Rs containing the  $\delta$  subunit have been shown to have an extra- or perisynaptic localization (40, 41) and give rise to tonic (sustained) GABA currents in neurons that express these GABA<sub>A</sub>R subunits (11). Recombinant GABA<sub>A</sub>Rs ( $\alpha 4/6\beta\delta$ ) known to mediate tonic currents, as well as  $\delta$  subunit-containing receptor-mediated tonic currents in neurons, are augmented by low alcohol concentrations reached during social alcohol consumption (9, 14, 42–44). Whereas  $\delta$  subunit-containing GABA<sub>A</sub>Rs comprise only between 5 and 10% of all of GABA<sub>A</sub>Rs in the mammalian brain (45), their constant or tonic activity (in marked contrast to synaptic receptors that open only for fractions of a second after synaptic GABA release) more than compensates in total





**Fig. 4.** A point mutation eliminates Ro15-4513-insensitive alcohol effects at high alcohol concentrations. (a) A single point mutation ( $\beta 3N265M$  in membrane-spanning segment TM2 of the  $\beta 3$  subunit) abolishes the Ro15-4513-resistant alcohol enhancement observed at high ethanol concentrations in  $\alpha 4\beta 3N265M\delta$  receptors. (b) Alcohol dose-response curve, determined by brief coapplications of ethanol and GABA  $EC_{50}$  (300 nM for  $\alpha 4\beta 3\delta$  and  $\alpha 4\beta 3N265M\delta$ , and 30  $\mu M$  GABA for  $\alpha 4\beta 3\gamma 2$  GABA $_A$ R). Currents through  $\alpha 4\beta 3N265M\delta$  GABA $_A$ R show a saturable alcohol enhancement and an  $EC_{50}$  of around 16 mM. The complete reversal of even very high dose alcohol effects by 300 nM Ro15-4513 in these experiments is surprising, because this behavior is not expected from an ideal competitive relationship of ligands with apparent dissociation constants (10 nM for Ro15-4513 and 16 mM for ethanol). The reasons why Ro15-4513 is so potent in antagonizing high-dose ethanol actions on functional receptors remain to be clarified.

charge transfer for the low abundance (46), and, therefore, extrasynaptic receptors are important regulators of neuronal excitability. The alcohol-induced augmentation of tonic currents in neurons and  $\delta$  subunit-containing GABA $_A$ Rs at relevant physiologic alcohol doses is expected to lead to a decrease in neuronal excitability in neurons expressing these subunits and make these receptors excellent candidates for mediating acute alcohol effects at intoxicating concentrations. Our demonstration that the behavioral alcohol antagonist Ro15-4513 leads to a dose-dependent inhibition of ethanol-induced current enhancement in recombinant  $\alpha 4\beta 3\delta$  GABA $_A$ Rs provides strong support for this notion.

**$\alpha 4\beta 3\delta$  GABA $_A$ R Are Sensitive to Certain BZs and BZ-Site Ligands.** The discovery that the BZ Ro15-4513 reduces alcohol actions on  $\alpha 4/6\beta 3\delta$  receptors is unexpected because it has been thought that the  $\gamma 2$  subunit is required for high-affinity binding of BZs and most BZ-site ligands (47). In recombinantly expressed functional receptors,  $\delta$  subunit-containing receptors have been shown to be insensitive to classical BZ agonists like diazepam and flunitrazepam (48, 49). Here, we reveal the activity of Ro15-4513 on  $\alpha 4/6\beta 3\delta$  receptors by their ability to act as an alcohol antagonist (Figs. 1, 2, and 4) whereas the binding of flumazenil and the BZ site ligand  $\beta$ -CCE is inferred from their ability to reverse Ro15-4513's alcohol antagonism (Fig. 2a). Consistent with the view that  $\alpha 4/6\beta 3\delta$  GABA $_A$ R are insensitive to classical BZ agonists, we show that flunitrazepam does not antagonize the alcohol antagonistic actions of Ro15-4513 on recombinant  $\alpha 4\beta 3\delta$  receptors (Fig. 2).

**Ethanol/Ro15-4513-Sensitive GABA $_A$ R in  $^{36}Cl^-$  Flux Assays Show Striking Similarities with Recombinantly Expressed  $\delta$  Subunit-Containing**

**GABA $_A$ R.** Our data on the Ro15-4513 reversal are in agreement with previous work that showed that Ro15-4513 can abolish ethanol augmentation of  $Cl^-$  flux in cerebral cortex synaptoneurosomal preparations and that the Ro15-4513's alcohol antagonism is reversed by flumazenil and  $\beta$ -CCE (17, 39). Several lines of evidence support the view that the ethanol-sensitive  $Cl^-$  flux through GABA $_A$ R in synaptosomal preparations may be mediated by alcohol-sensitive extrasynaptic receptors: (i) like  $\delta$  subunit-containing receptors, this  $Cl^-$  flux seems highly sensitive to GABA and muscimol (17, 29), and the ability to carry sustained  $^{36}Cl^-$  flux suggests that they show slow desensitization; (ii) both  $\delta$  subunit-containing receptors, as well as the  $^{36}Cl^-$  flux, are strikingly similar in their low-dose response to physiologically relevant (3–30 mM) ethanol concentrations; (iii) we show here with recombinant receptors that, as previously observed in the  $^{36}Cl^-$  flux assays (17), 30 mM ethanol augmentation is completely reversed by 300 nM of the behavioral alcohol antagonist Ro15-4513.

**Why Are GABA $_A$ R as Alcohol Targets Controversial?** Our findings also provide an explanation for the controversial findings concerning the alcohol sensitivity of GABA $_A$ Rs in native neurons. Whereas most synaptic physiologists failed to find evidence for low-dose alcohol effects on GABAergic synaptic transmission (42, 50), there is abundant evidence that many neurons can respond to physiological (3–30 mM) ethanol concentrations at conditions that favor the detection of highly GABA-sensitive nondesensitizing extrasynaptic GABA $_A$ Rs (6, 51). Such conditions are, for example, the prolonged application of relatively low GABA and ethanol concentrations, like those resulting from local application of GABA/ethanol in the vicinity of neurons, conditions that synaptic physiologists might have considered nonphysiological. Consistent with this view and the results presented here, it has been shown that such low-dose alcohol enhancement of GABA currents in neurons (51) and the (presumably) resulting changes in firing frequency (52) can be reversed by the alcohol antagonist Ro15-4513 (51, 52). The fact that not all neurons express these alcohol/Ro15-4513-sensitive  $\delta$  subunit-containing GABA $_A$ Rs is consistent with the observation that not all neurons are sensitive to pharmacologically relevant ethanol concentrations (6, 42, 51). Another complicating factor is that electrophysiological studies are often performed on neurons and slices harvested from immature brains that do not yet express  $\delta$  subunit-containing extrasynaptic GABA $_A$ R (53). Consistent with a slow onset of expression of  $\delta$  subunit-containing receptors during development and adolescence in rodents, reaching mature levels at around sexual maturation (54, 55), is the finding that younger animals are less sensitive to the sedative and motor-impairing effects of ethanol (56).

Whereas our experimental data explain and extend >20 yr of often puzzling observations concerning ethanol actions on GABA $_A$ Rs, a recent report that fails to reproduce low-dose ethanol enhancement on  $\alpha 4\beta 3\delta$  receptors, in particular negative results from the attempted expression of human  $\alpha 4\beta 3\delta$  receptors in oocytes and mammalian cell lines (57), likely ensures that our findings and related work (e.g., refs. 14, 17, 29, and 39) will remain controversial. We are puzzled by the differences, not only in the apparent lack of alcohol sensitivity, but also in the peak currents published by Borghese *et al.* (57), which are five times higher than we see with the (rat) clones that we provided to them. In marked contrast, the GABA responses from the human clones expressed in oocytes reported by Borghese *et al.* (57) are marginal, indicating a possible lack of expression of one or more of their human clones.

In our hands, variable ethanol responses in  $\alpha 4\beta 3\delta$  are likely due to the tendency of oocytes to express receptors that are composed of  $\alpha\beta$  subunits alone. GABA $_A$ Rs composed of  $\alpha\beta$  alone are insensitive to relevant alcohol concentrations and to classical BZs but show high sensitivity to etomidate, propofol, and steroid anesthetics (58, 59). The phenomenon of  $\alpha\beta$

GABA<sub>A</sub>R expression in oocytes has been well documented for  $\gamma 2$  subunit-containing receptors and explains the observed variability in BZ effects in  $\alpha 1\beta 2\gamma 2$  receptors expressed in oocytes (60). The expression of  $\alpha\beta$  receptors in oocytes, highly sensitive to GABA<sub>A</sub>R anesthetics, is also the likely explanation for a previous challenge (58) to the finding that  $\epsilon$  subunit-containing GABA<sub>A</sub>Rs lack anesthetic enhancement (59).

**Behavioral Alcohol Effects Antagonized by Ro15-4513.** The alcohol antagonist Ro15-4513 has been reported to prevent many acute alcohol effects. These effects include increased exploration and locomotion at very low doses (0.25, 0.5, and 0.75 g/kg in rats) (61), the anxiolytic effects at low doses (1 g/kg) (17, 39), and sedative, motor-impairing as well as the amnesic effects at moderate alcohol doses (2 g/kg) (17, 19, 21, 62). In addition, the observation that Ro15-4513 reduces alcohol self-administration (23, 24, 63) suggests that the rewarding effects of ethanol might be mediated by ethanol/Ro15-4513-sensitive GABA<sub>A</sub>Rs. However, Ro15-4513 does not prevent all ethanol effects: at higher alcohol doses ( $\geq 2$  g/kg in rats), Ro15-4513 significantly reduces, but does not prevent the anesthetic ("sleep"-inducing) effects of ethanol, and Ro15-4513 does not prevent the hypothermic effects of ethanol (32, 64). In addition, Ro15-4513 does not prevent lethal effects at massive alcohol doses (20, 27). This finding suggests that, at high alcohol doses, Ro15-4513-insensitive ethanol targets mediate the anesthetic ethanol effects. Our data on recombinant  $\alpha 4\beta 3\delta$  GABA<sub>A</sub>Rs suggest the possibility that, at such high alcohol concentrations (blood alcohol levels  $> 30$  mM), Ro15-4513-sensitive recombinant  $\alpha 4/6\beta 3\delta$  receptors have a high ethanol ( $> 80\%$ ) occupancy and are therefore close to saturated.

Established targets of Ro15-4513-insensitive alcohol actions are alcohol-sensitive G protein-gated (GIRK) potassium channels (33) that likely contribute to the analgesic (4) and hypothermic effects of ethanol (34). Other potential alcohol targets that might contribute to Ro15-4513-insensitive acute alcohol actions are NMDA (1), glycine (2), and adenosine receptors (65), and voltage- and Ca<sup>2+</sup>-activated (BK) potassium channels (66, 67). In addition, we show here that the Ro15-4513-insensitive alcohol action on  $\alpha 4\beta 3\delta$  GABA<sub>A</sub>R is abolished by a mutation ( $\beta 3N265M$ ) in the  $\beta 3$  subunit. It is therefore possible that GABA<sub>A</sub>Rs (including the abundant synaptic  $\gamma 2$ -containing GABA<sub>A</sub>Rs) may contribute to high-dose ( $\geq 30$  mM) Ro15-4513-resistant, anesthetic ethanol actions.

**Potential Mechanisms of Alcohol Antagonism by Ro15-4513.** A comparison of the structure of the alcohol and BZ antagonist Ro15-4513 and flumazenil, which differ only at a single moiety, suggests a possible mechanism of alcohol antagonism. The larger azido group (at the C7 position of the BZ ring) might be the group that occupies the alcohol-binding site on the receptor. Flumazenil and the  $\beta$ -carbolines  $\beta$ -CCE and FG7142 likely antagonize Ro15-4513 alcohol antagonist actions by displacing Ro15-4513 from its binding site. Flumazenil,  $\beta$ -CCE, and FG7142 do not act as alcohol antagonists by themselves, because they might fit together with ethanol in the Ro15-4513-binding pocket. Therefore, we think that the most parsimonious explanation for Ro15-4513's alcohol antagonism is that the unique azido group in Ro15-4513 occupies the alcohol-binding site. However, there could be other possible mechanisms: e.g., the azido group in Ro15-4513 may cause allosteric changes in these receptors. We have used native immunopurified and recombinant expressed  $\delta$  subunit-containing receptors in [<sup>3</sup>H]Ro15-4513-binding assays to show further evidence that Ro15-4513 and ethanol have a competitive relationship on  $\alpha 4/6\beta 3\delta$  receptors (68).

**Alcohol Receptors as Potential Drug Targets.** Our finding that alcohol effects on  $\alpha 4\beta 3\delta$  receptors can be reversed by the BZ Ro15-4513

and that this action in turn is sensitive to flumazenil and the BZ site ligand  $\beta$ -CCE suggests that these receptors likely contain a modified BZ-binding site. Such previously unrecognized BZ sites, on less abundant but functionally important GABA<sub>A</sub>R subtypes, provide opportunity for new drug development. For example, it might be advantageous to synthesize alcohol antagonists that lack the inverse agonist effects of Ro15-4513 on certain GABA receptor subtypes. In addition, given Ro15-4513's structural similarity with the general BZ antagonist flumazenil, it is not surprising that Ro15-4513 is also a general BZ antagonist, and like flumazenil likely has a fairly short half-life ( $\approx 30$  min) *in vivo*. Furthermore, the alcohol antagonist Ro15-4513 and the clinically used general BZ antagonist flumazenil are highly hydrophobic with likely poor bioavailability when applied orally or i.p. (flumazenil is administered intravenously in the clinic as a BZ antidote). Given the high incidence of ethanol intoxication cases in emergency rooms and the short half-life of flumazenil, leading to re sedation in cases where it is used to antagonize much longer acting BZs, the identification of  $\alpha 4/6\beta 3\delta$  GABA<sub>A</sub>Rs as ethanol/Ro15-4513 targets may spur the development of combined BZ/alcohol antagonists with longer half-life. Such combined antagonists, without inverse agonist activity and better solubility, not only might be useful in the clinic, but also may provide research tools for a precise dissection of the contribution of GABA<sub>A</sub>R subtypes to certain aspects of acute ethanol actions.

We show that the BZ-site ligand  $\beta$ -CCE, a  $\beta$ -carboline, is an agonist on  $\alpha 4\beta 3\delta$  receptors and like EtOH allosterically enhances the GABA response without directly activating these receptors. The activation of  $\alpha 4\beta 3\delta$  GABA<sub>A</sub>Rs by  $\beta$ -CCE provides proof of principle that it might be possible to develop specific alcohol receptor agonists. Ideally, such positive modulators (alcohol mimetics) should be specific for the EtOH/Ro15-4513 site and lack activity (like the inverse agonist activity of  $\beta$ -CCE) on classical GABA<sub>A</sub>R  $\alpha 1$ , -2, -3, and -5 $\gamma 2$  BZ sites. Such specific alcohol receptor agonists could be useful to harness ethanol receptors for therapeutic purposes by mimicking the anxiolytic, anti-depressive, and mood-enhancing actions of alcohol, without the undesired effects like liver toxicity and other toxic effects of metabolites like acetaldehyde.

## Materials and Methods

**Electrophysiology.** Clones used were as described previously and were confirmed by sequencing to ensure that they were free of errors and agreed with the consensus sequences for rat  $\alpha 4$ ,  $\alpha 6$ ,  $\beta 3$ , and  $\delta$  subunit proteins (9). Mutagenesis was performed by using the QuikChange site-directed mutagenesis kit (Stratagene). cRNA was transcribed after plasmid linearization by using the mMessage mMachine kit (Ambion, Austin, TX). Transcripts were purified by LiCl precipitation, and RNA concentration was determined on a gel and by photometry. Oocytes were coinjected with a mixture of  $\alpha$ ,  $\beta$ , and  $\delta$  (or  $\gamma 2$ ) subunits in a 1:1:5 (or 1:1:10) subunit molar ratio. Currents were measured at room temperature (22°C–24°C) in the two-electrode voltage clamp configuration at  $-80$  mV holding potential with an Axoclamp 2A Axon Instruments (Union City, CA) amplifier. Two electrode voltage clamp on *Xenopus* oocytes was performed in ND96 salt solution (96 mM NaCl/2 mM KCl/1.8 mM CaCl<sub>2</sub>/1 mM MgCl<sub>2</sub>/5 mM Hepes, pH 7.2). Because of the slow onset in the expression of highly alcohol-sensitive  $\delta$  subunit-containing receptors, oocytes were measured 7–14 days after injection. Currents were measured either in a tonic current mimicking mode as steady-state current (Figs. 1 and 2), or by brief ( $< 10$  s) coapplications of 300 nM GABA with drugs (ethanol, BZ-site ligands), to evoke peak currents, followed by a recovery time of at least 1 min.

**Reagents.** Hoffman-LaRoche (Nutley, NJ) kindly provided Ro15-4513, flumazenil or Ro15-1788 (ethyl 8-fluoro-5,6-dihydro-5-methyl-6-oxo-4H-imidazo[1,5-a][1,4]benzodiazepine-3-carboxylate), diazepam, and flunitrazepam. DMCM was a gift



from Ferrosan (Copenhagen), and FG7142 (*N*-methyl- $\beta$ -carboline-3-carboxamide) and  $\beta$ -CCE were provided by Schering. Ethanol, GABA, and bicuculline were purchased from Sigma. Compounds were dissolved in DMSO as a 10-mM stock solution and were used at the indicated concentrations. DMSO at final concentrations used did not lead to changes in GABA receptor currents (data not shown).

- Popp, R. L., Lickteig, R. L. & Lovinger, D. M. (1999) *J. Pharmacol. Exp. Ther.* **289**, 1564–1574.
- Davies, D. L., Trudell, J. R., Mihic, S. J., Crawford, D. K. & Alkana, R. L. (2003) *Alcohol Clin. Exp. Res.* **27**, 743–755.
- Ikeda, K., Kobayashi, T., Kumanishi, T., Yano, R., Sora, I. & Niki, H. (2002) *Neurosci. Res.* **44**, 121–131.
- Blednov, Y. A., Stoffel, M., Alva, H. & Harris, R. A. (2003) *Proc. Natl. Acad. Sci. USA* **100**, 277–282.
- Liljequist, S. & Engel, J. (1982) *Psychopharmacology* **78**, 71–75.
- Aguayo, L. G., Peoples, R. W., Yeh, H. H. & Yevenes, G. E. (2002) *Curr. Top. Med. Chem.* **2**, 869–885.
- Hu, W. Y., Reiffenstein, R. J. & Wong, L. (1987) *Alcohol Drug Res.* **7**, 107–117.
- Khanna, J. M., Kalant, H., Chau, A. & Shah, G. (1998) *Pharmacol. Biochem. Behav.* **59**, 511–519.
- Wallner, M., Hanchar, H. J. & Olsen, R. W. (2003) *Proc. Natl. Acad. Sci. USA* **100**, 15218–15223.
- Mihalek, R. M., Bowers, B. J., Wehner, J. M., Kralic, J. E., VanDoren, M. J., Morrow, A. L. & Homanics, G. E. (2001) *Alcohol Clin. Exp. Res.* **25**, 1708–1718.
- Farrant, M. & Nusser, Z. (2005) *Nat. Rev. Neurosci.* **6**, 215–229.
- Benson, J. A., Low, K., Keist, R., Möhler, H. & Rudolph, U. (1998) *FEBS Lett.* **431**, 400–404.
- Korpi, E. R., Kleingoor, C., Kettenmann, H. & Seeburg, P. H. (1993) *Nature* **361**, 356–359.
- Hanchar, H. J., Dodson, P. D., Olsen, R. W., Otis, T. S. & Wallner, M. (2005) *Nat. Neurosci.* **8**, 339–345.
- Bonetti, E. P., Burkhard, W. P., Gabl, M. & Möhler, H. (1985) *Br. J. Pharmacol.* **86**, 463P.
- Polc, P. (1985) *Br. J. Pharmacol.* **86**, 465P.
- Suzdak, P. D., Glowa, J. R., Crawley, J. N., Schwartz, R. D., Skolnick, P. & Paul, S. M. (1986) *Science* **234**, 1243–1247.
- Syapin, P. J., Jones, B. L., Kobayashi, L. S., Finn, D. A. & Alkana, R. L. (1990) *Brain Res. Bull.* **24**, 705–709.
- Bonetti, E. P., Burkard, W. P., Gabl, M., Hunkeler, W., Lorez, H. P., Martin, J. R., Möhler, H., Osterrieder, W., Pieri, L. & Polc, P. (1988) *Pharmacol. Biochem. Behav.* **31**, 733–749.
- Kolata, G. (1986) *Science* **234**, 1198–1199.
- Nabeshima, T., Tohyama, K. & Kameyama, T. (1988) *Eur. J. Pharmacol.* **155**, 211–217.
- June, H. L., Hughes, R. W., Spurlock, H. L. & Lewis, M. J. (1994) *Psychopharmacology* **115**, 332–339.
- Petry, N. M. (1995) *Psychopharmacology* **121**, 192–203.
- Rassnick, S., D'Amico, E., Riley, E. & Koob, G. F. (1993) *Alcohol Clin. Exp. Res.* **17**, 124–130.
- Britton, K. T., Ehlers, C. L. & Koob, G. F. (1988) *Science* **239**, 648–650.
- Suzdak, P. D., Glowa, J. R., Crawley, J., Skolnick, P. & Paul, S. M. (1987) *Science* **239**, 649–650.
- Lister, R. G. & Nutt, D. J. (1987) *Trends Neurosci.* **10**, 223–225.
- Lister, R. G. & Nutt, D. J. (1988) *Pharmacol. Biochem. Behav.* **31**, 731.
- Suzdak, P. D., Schwartz, R. D., Skolnick, P. & Paul, S. M. (1986) *Proc. Natl. Acad. Sci. USA* **83**, 4071–4075.
- Allan, A. M. & Harris, R. A. (1986) *Life Sci.* **39**, 2005–2015.
- Suzdak, P. D., Paul, S. M. & Crawley, J. N. (1988) *J. Pharmacol. Exp. Ther.* **245**, 880–886.
- Syapin, P. J., Gee, K. W. & Alkana, R. L. (1987) *Brain Res. Bull.* **19**, 603–605.
- Kobayashi, T., Ikeda, K., Kojima, H., Niki, H., Yano, R., Yoshioka, T. & Kumanishi, T. (1999) *Nat. Neurosci.* **2**, 1091–1097.
- Costa, A. C., Stasko, M. R., Stoffel, M. & Scott-McKean, J. J. (2005) *J. Neurosci.* **25**, 7801–7804.
- Jurd, R., Arras, M., Lambert, S., Drexler, B., Siegwart, R., Crestani, F., Zaugg, M., Vogt, K. E., Ledermann, B., Antkowiak, B. & Rudolph, U. (2003) *FASEB J.* **17**, 250–252.
- Belelli, D., Lambert, J. J., Peters, J. A., Wafford, K. & Whiting, P. J. (1997) *Proc. Natl. Acad. Sci. USA* **94**, 11031–11036.
- Wafford, K. A., Bain, C. J., Quirk, K., McKernan, R. M., Wingrove, P. B., Whiting, P. J. & Kemp, J. A. (1994) *Neuron* **12**, 775–782.
- Mihic, S. J., Ye, Q., Wick, M. J., Koltchine, V. V., Krasowski, M. D., Finn, S. E., Mascia, M. P., Valenzuela, C. F., Hanson, K. K., Greenblatt, E. P., et al. (1997) *Nature* **389**, 385–389.
- Glowa, J. R., Crawley, J., Suzdak, P. D. & Paul, S. M. (1988) *Pharmacol. Biochem. Behav.* **31**, 767–772.
- Nusser, Z., Sieghart, W. & Somogyi, P. (1998) *J. Neurosci.* **18**, 1693–1703.
- Wei, W., Zhang, N., Peng, Z., Houser, C. R. & Mody, I. (2003) *J. Neurosci.* **23**, 10650–10661.
- Wei, W., Faria, L. C. & Mody, I. (2004) *J. Neurosci.* **24**, 8379–8382.
- Carta, M., Mameli, M. & Valenzuela, C. F. (2004) *J. Neurosci.* **24**, 3746–3751.
- Liang, J., Zhang, N., Cagetti, E., Houser, C. R., Olsen, R. W. & Spigelman, I. (2006) *J. Neurosci.* **26**, 1749–1758.
- Whiting, P., Wafford, K. & McKernan, R. M. (2000) in *GABA in the Nervous System: The View at Fifty Years*, eds Martin, D. L. & Olsen, R. W. (Lippincott Williams & Wilkins, Philadelphia), pp. 113–126.
- Nusser, Z. & Mody, I. (2002) *J. Neurophysiol.* **87**, 2624–2628.
- Sigel, E. (2002) *Curr. Top. Med. Chem.* **2**, 833–839.
- Saxena, N. C. & Macdonald, R. L. (1996) *Mol. Pharmacol.* **49**, 567–579.
- Brown, N., Kerby, J., Bonnert, T. P., Whiting, P. J. & Wafford, K. A. (2002) *Br. J. Pharmacol.* **136**, 965–974.
- Criswell, H. E. & Breese, G. R. (2005) *Neuropsychopharmacology* **30**, 1407–1425.
- Reynolds, J. N., Prasad, A. & MacDonald, J. F. (1992) *Eur. J. Pharmacol.* **224**, 173–181.
- Palmer, M. R., van Horne, C. G., Harlan, J. T. & Moore, E. A. (1988) *J. Pharmacol. Exp. Ther.* **247**, 1018–1024.
- Ming, Z., Knapp, D. J., Mueller, R. A., Breese, G. R. & Criswell, H. E. (2001) *Brain Res.* **920**, 117–124.
- Laurie, D. J., Wisden, W. & Seeburg, P. H. (1992) *J. Neurosci.* **12**, 4151–4172.
- Brickley, S. G., Revilla, V., Cull-Candy, S. G., Wisden, W. & Farrant, M. (2001) *Nature* **409**, 88–92.
- White, A. M., Truesdale, M. C., Bae, J. G., Ahmad, S., Wilson, W. A., Best, P. J. & Swartzwelder, H. S. (2002) *Pharmacol. Biochem. Behav.* **73**, 673–677.
- Borghese, C. M., Sturustovu, S. I., Ebert, B., Herd, M. B., Belelli, D., Lambert, J. J., Marshall, G., Wafford, K. A. & Harris, R. A. (2006) *J. Pharmacol. Exp. Ther.* **316**, 1360–1368.
- Thompson, S. A., Bonnert, T. P., Cagetti, E., Whiting, P. J. & Wafford, K. A. (2002) *Neuropharmacology* **43**, 662–668.
- Davies, P. A., Hanna, M. C., Hales, T. G. & Kirkness, E. F. (1997) *Nature* **385**, 820–823.
- Boileau, A. J., Baur, R., Sharkey, L. M., Sigel, E. & Czajkowski, C. (2002) *Neuropharmacology* **43**, 695–700.
- June, H. L. & Lewis, M. J. (1994) *Psychopharmacology* **116**, 309–316.
- Dar, M. S. (1995) *Pharmacol. Biochem. Behav.* **52**, 217–223.
- June, H. L., Lummis, G. H., Colker, R. E., Moore, T. O. & Lewis, M. J. (1991) *Alcohol Clin. Exp. Res.* **15**, 406–411.
- Hoffman, P. L., Tabakoff, B., Szabo, G., Suzdak, P. D. & Paul, S. M. (1987) *Life Sci.* **41**, 611–619.
- Dar, M. S., Mustafa, S. J. & Wooles, W. R. (1983) *Life Sci.* **33**, 1363–1374.
- Davies, A. G., Pierce-Shimomura, J. T., Kim, H., VanHoven, M. K., Thiele, T. R., Bonci, A., Bargmann, C. I. & McIntire, S. L. (2003) *Cell* **115**, 655–666.
- Liu, P., Xi, Q., Ahmed, A., Jaggart, J. H. & Dopico, A. M. (2004) *Proc. Natl. Acad. Sci. USA* **101**, 18217–18222.
- Hanchar, H. J., Chutsrinopkun, P., Meera, P., Supavilai, P., Sieghart, W., Wallner, M. & Olsen, R. W. (2006) *Proc. Natl. Acad. Sci. USA* **103**, 8546–8551.