

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/305077017>

Recent Advances in the Rational Design and Optimization of Antibacterial Agents

Article in *Medicinal Chemistry Communication* · September 2016

DOI: 10.1039/C6MD00232C

CITATION

1

READS

109

4 authors, including:



Jesse Jones

The University of Tennessee Health Science C...

2 PUBLICATIONS 1 CITATION

SEE PROFILE



Kristopher Virga

Presbyterian College

15 PUBLICATIONS 334 CITATIONS

SEE PROFILE



Kirk Hevener

University of Tennessee, Health Science Center

25 PUBLICATIONS 436 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Investigation of Streptococcal Topoisomerase I as a Narrow-Spectrum Antibacterial Target [View project](#)



Naegleria fowleri [View project](#)

All content following this page was uploaded by [Kirk Hevener](#) on 18 July 2016.

The user has requested enhancement of the downloaded file. All in-text references [underlined in blue](#) are added to the original document and are linked to publications on ResearchGate, letting you access and read them immediately.

CrossMark
click for updates

Cite this: DOI: 10.1039/c6md00232c

Recent advances in the rational design and optimization of antibacterial agents†

Jesse A. Jones,^a Kristopher G. Virga,^b Giuseppe Gumina^b and Kirk E. Hevener^{*a}

This review discusses next-generation antibacterial agents developed using rational, or targeted, drug design strategies. The focus of this review is on small-molecule compounds that have been designed to bypass developing bacterial resistance, improve the antibacterial spectrum of activity, and/or to optimize other properties, including physicochemical and pharmacokinetic properties. Agents are discussed that affect known antibacterial targets, such as the bacterial ribosome, nucleic acid binding proteins, and proteins involved in cell-wall biosynthesis; as well as some affecting novel bacterial targets which do not have currently marketed agents. The discussion of the agents focuses on the rational design strategies employed and the synthetic medicinal chemistry and structure-based design techniques utilized by the scientists involved in the discoveries, including such methods as ligand- and structure-based strategies, structure-activity relationship (SAR) expansion strategies, and novel synthetic organic chemistry methods. As such, the discussion is limited to small-molecule therapeutics that have confirmed macromolecular targets and encompasses only a fraction of all antibacterial agents recently approved or in late-stage clinical trials. The antibacterial agents selected have been recently approved for use on the U.S. or European markets or have shown promising results in phase 2 or phase 3 U.S. clinical trials.

Received 26th April 2016,
Accepted 5th July 2016

DOI: 10.1039/c6md00232c

www.rsc.org/medchemcomm

Introduction

Beginning with the discovery of penicillin in 1928, the early twentieth century ushered in an unprecedented “golden era” of antibacterial drug discovery. Phenotypic whole-cell screening of natural products predominated this era and carried with it a success that is yet to be reproduced. Effective drug discovery and development later transitioned into the mid- to

^a Department of Biomedical and Pharmaceutical Sciences, Idaho State University, 1311 E. Central Drive, Meridian, ID 83642-7991, USA.

E-mail: khevener@pharmacy.isu.edu; Fax: +1 312 413 9303; Tel: +1 208 373 1829

^b Department of Pharmaceutical Sciences, Presbyterian College School of Pharmacy, 307 North Broad Street, Clinton, SC 29325, USA

† The authors declare no competing interests.

**Jesse A. Jones**

Jesse A. Jones, Pharm.D., received his B.S. from Boise State University in 2008. He then started work toward a Pharm.D. in Pocatello, ID at Idaho State University (ISU), which he completed in Meridian, ID at the ISU Meridian Health Science Center in 2012. He is currently a Ph.D. student in Pharmaceutical Sciences with a concentration in Drug Discovery at the ISU Meridian Health Science Center under Dr. Kirk E. Hevener, where he

has been for three years. His research interests include the rational discovery and design of antimicrobial agents, including the use of molecular biochemistry and structural biology.

**Kristopher Virga**

Kristopher Virga, Ph.D., is an Associate Professor within the Department of Pharmaceutical and Administrative Sciences at the Presbyterian College, School of Pharmacy. Dr. Virga has ten years of teaching and research experience as a faculty member within a college of pharmacy. Dr. Virga completed his Ph.D. at University of Tennessee Health Science Center and a postdoctoral fellowship at University of Notre Dame. His disciplinary

area of expertise is in medicinal chemistry. His main research focus is on the development of novel antimicrobial agents with a particular interest in targeting bacterial biofilms.

late-twentieth century in the form of a strong “medicinal chemistry era.” This period was mainly oriented around synthetic modifications of known compounds, high-throughput screening (HTS), and library design.¹ It was initially hoped that this subsequent age of antibacterial discovery and development would be a triumph over the long reign of bacterial infections, but instead it has more recently proven to be a simple prelude to the modern antibacterial era—an age marked by the looming threat of expanding bacterial resistance coupled with the stark reality of an inversely diminishing antibacterial pipeline.^{2–4} As opposed to those bygone golden and medicinal chemistry eras, the contemporary “resistance era” is marked by a drug discovery climate that is comparatively less fruitful and geared around more time-consuming rationalized design modalities featuring target-based HTS campaigns that make use of commonly exhaustive tools like combinatorial libraries.⁵

Continuing along the antibacterial timeline, the commonly forecasted “post-antibiotic era” showcases a bleak future in which it is estimated that by the year 2050 the number of human deaths attributed to resistant microorganisms will surpass 10 million per year, up from the current 700 000 per year.² To combat this issue, a multi-faceted approach has commenced. The U.S. has recently launched infectious disease initiatives including legislation such as the Generating Antibiotics Incentives Now Act (GAIN Act) aimed at fast-tracking Qualified Infectious Disease Products (QIDPs). Europe has also responded with its own countermeasures, such as Innovative Medicines Initiative’s (IMI’s) ENABLE project under the “New Drugs 4 Bad Bugs” (ND4BB) program aimed at battling Gram-negative bacteria.

To meet the antibacterial drug discovery and development challenges at hand, modern research efforts have further expanded upon *rational* design and optimization strategies—such efforts addressing specific targets or problems, such as cell penetration.⁵ These efforts revolve around improved ac-

tivity, physicochemical properties, pharmacokinetics, adverse effects, off-target activities, and so forth. Moreover, researchers have recently appeared to “re-discover” natural products chemistry, additionally employing novel approaches like modern semi-synthetic techniques.^{6,7} With newly implemented incentivization and modern advances in research methods, it is hoped that the threat of antibacterial resistance will be met with appropriate countermeasures. It is therefore the objective of this review to discuss the latest progress and developments in the rational design and optimization of antibacterial agents recently approved or in late-phase trials.

Discussion

Advanced generation cephalosporins

Recent advances in the chemistry of the β -lactam antibiotics have resulted in the approval or near approval of several advanced (or 5th) generation cephalosporins with enhanced spectra of activity, including methicillin-resistant *Staphylococcus aureus* (MRSA), and *Pseudomonas aeruginosa* (PA). While the general structure–activity relationship (SAR) of the cephalosporins has been covered well in previous communications,⁸ there has been some notable medicinal chemistry applied to the advanced generation cephalosporins that merits discussion here. The classical cephalosporin scaffold (Fig. 1) includes a four-membered β -lactam ring fused with a dihydrothiazine ring (cephem) that contains a double bond between the 2- and 3-position carbons, and a 2-position carboxylate. Essential to the activity of the cephalosporin antibiotics is a functional group substitution at the 3-position that can stabilize a negative charge resulting from β -lactam ring opening and increase the overall reactivity of the β -lactam ring when interacting with target transpeptidase enzymes.

Ceftaroline (Teflaro® in the U.S., Zinforo® in Europe, Forest Laboratories) was approved for use in the U.S. for treatment



Giuseppe Gumina

Giuseppe Gumina, Ph.D. is Associate Professor of Pharmaceutical and Administrative Sciences at Presbyterian College, School of Pharmacy. Prior to his current position, he held academic positions at South University School of Pharmacy and the Medical University of South Carolina, College of Pharmacy. His postgraduate research in medicinal chemistry was conducted at the University of Georgia. His undergraduate and graduate studies

in medicinal chemistry with emphasis in synthetic organic chemistry were completed at the University of Catania, Italy. His research focuses on the design and synthesis of novel nucleoside analogs as antimicrobial and anticancer agents.



Kirk E. Hevener

Kirk E. Hevener, Pharm.D., Ph.D., is Assistant Professor of Biomedical and Pharmaceutical Sciences at Idaho State University, College of Pharmacy. Dr. Hevener earned a BS in Organic Chemistry from Tennessee State University followed by a Pharm.D. and Ph.D. from the University of Tennessee Health Science Center. His research expertise is in medicinal chemistry, computational chemistry, and structural biology. Before coming to

Idaho State University, Dr. Hevener completed postdoctoral fellowships at St. Jude Children’s Research Hospital and the University of Illinois at Chicago.




Fig. 1 Cephem ring, structure and positional numbering.

of community-acquired bacterial pneumonia (CABP) and acute bacterial skin and skin structure infections (ABSSSI) in 2010. Ceftaroline is marketed as the prodrug Ceftaroline fosamil, which incorporates a hydrolysable phosphono group to improve aqueous solubility (Fig. 2A).⁹ The drug has excellent Gram positive coverage, including activity against resistant strains of *Streptococcus pneumoniae* and MRSA, and moderate Gram negative coverage, but is less active against *Pseudomonas aeruginosa*, enterococci, or clinically relevant anaerobic organisms.^{10–12} Key to the drug's activity is a high binding affinity for the penicillin-binding protein 2a (PBP2a), which is known to play a significant role in resistance to anti-staphylococcal β -lactams, and PBP2x, found in β -lactam-resistant *S. pneumoniae*. As can be seen from the chemical structure (Fig. 2A), ceftaroline retains an oxime group as part of the 7-position substituent, in this case an ethyl oxime. This is also seen in the third- and fourth-generation cephalosporins, and is known to confer resistance to some β -lactamase enzymes. The 1,2,4-thiadiazine ring component of the 7-position substituent is believed to

improve the drug's penetration into Gram-negative organisms, as well as contribute to overall target affinity.¹³ The 1,3-thiazole ring, connected to the cephem ring at the 3-position by a unique sulfide bond, has been reported to play a key role in the improved affinity of ceftaroline for PBP2a and the resulting anti-MRSA activity.¹³ Finally, the quaternary nitrogen of the pyridine ring in the 3-position substituent contributes a positive charge to the overall zwitterion ceftaroline and improves drug penetration into Gram-negative organisms.

Ceftobiprole (Zevtera®, Basilea Pharmaceutica) was approved in Europe and Canada for the treatment of community-acquired pneumonia (CAP) and hospital-acquired pneumonia (HAP), excluding ventilator-associated pneumonia (VAP) in 2013. After initially being postponed in Europe and discontinued in Canada in 2010 due to recommendations from regulatory agencies in Europe and the U.S. regarding clinical trial concerns, Ceftobiprole is now marketed as ceftobiprole medocartil sodium, a prodrug incorporating a solubilizing group attached to the pyrrolidine nitrogen (Fig. 2B) by a hydrolysable carbamate ester.^{14,15} The drug has reported broad-spectrum activity against many Gram positive and Gram negative organisms, including resistant strains of *S. aureus* (MRSA & VRSA), *S. pneumoniae* (PRSP & CRSP), *P. aeruginosa*, and *Enterobacteriaceae*.^{16,17} It is less active against clinically relevant anaerobic organisms, including *Prevotella* species and *B. fragilis* group species.¹⁸ The presence of the oxime functionality in the 7-position




Fig. 2 Advanced generation cephalosporins recently approved or in clinical trials. A. Ceftaroline fosamil. B. Ceftobiprole medocartil (prodrug moiety in red). C. Ceftolozane. D. S-649266.

substituent confers a degree of β -lactamase resistance to ceftobiprole similar to that seen in the 3rd and 4th generation cephalosporins, however the drug is not resistant to the extended-spectrum β -lactamases (ESBL) produced by resistant strains of *Enterobacteriaceae*.¹⁹ Similar to ceftaroline, the activity of ceftobiprole is attributed to an increased affinity for PBP2a (*S. aureus*) and PBP2x (*S. pneumoniae*). The co-crystal structure of the complex of ceftobiprole bound to PBP2a has provided key insights in the nature of the binding interactions of the drug to its target (PDB accession code 4DKI).²⁰ Key to the increased binding affinity of ceftobiprole for PBP2a appears to be the vinyl-pyrrolidinone component of the 3-position substituent, which interacts with a narrow groove of the PBP2a active site, forming π -stacking interactions with a nearby tyrosine residue (Tyr-446) and general hydrophobic interactions with other nearby residues, resulting in a favorable positioning of the β -lactam ring for acylation by the key serine residue (Ser-403). Binding-induced conformational changes in secondary structure elements near the active site also appear to facilitate the acylation reaction.²⁰

Another advanced generation cephalosporin, ceftolozane, was approved in the U.S. for use in combination with the β -lactamase inhibitor, tazobactam, for treatment of complicated intra-abdominal infections (in combination with metronidazole) and complicated urinary tract infections in 2014 (ZerbaxaTM, Merck). Ceftolozane lacks activity against many clinically relevant Gram positive organisms, with no appreciable activity against MRSA. However, the addition of tazobactam gives the combination product excellent Gram negative activity including potent activity against many resistant *P. aeruginosa* strains and most ESBL-producing *Enterobacteriaceae*.^{21,22} The combination (ceftolozane/tazobactam) shows highly variable activity against anaerobic organisms, particularly against *B. fragilis* species, which necessitates the use in combination with metronidazole for complicated intra-abdominal infections (cIAIs).²³ Structurally, ceftolozane possesses the 7-substituent oxime functionality conferring a degree of β -lactamase resistance similar to the 3rd and 4th generation cephalosporins. Similar to ceftazidime, the oxime functionality incorporates a dimethyl acetic acid moiety that is believed to enhance activity against *P. aeruginosa* (Fig. 2C).²⁴ At the 3-position, the bulky, substituted pyrazole ring prevents hydrolysis of the β -lactam ring by steric hindrance and confers some stability to AmpC β -lactamases commonly seen in *P. aeruginosa*.^{25,26} Unlike ceftaroline and ceftobiprole, the 3-position substituent of ceftolozane does not appear to improve the drug's affinity for PBP2a, which explains the compounds lack of activity against MRSA, and the drug has low overall affinity for PBP4, similar to ceftazidime, which explains its very weak induction of AmpC expression.²⁷ However, the ceftolozane has particularly strong affinity for PBP3, at least 2-fold higher than ceftazidime, which explains its potent *P. aeruginosa* activity.²⁷ Indeed, the 3-substituent pyrazole substituents were intentionally designed to maximize this affinity and hence activity against *P. aeruginosa*.^{24,26}

Compound S-649266 (Shionogi, Inc.) is an unapproved, advanced generation cephalosporin currently in phase 2 clinical trials in the U.S for treatment of complicated urinary tract infections (cUTI). The compound is notable for its activity against bacterial strains producing 'K. pneumoniae Carbapenemase' (KPC) and 'New Delhi Metallo- β -lactamase' (NDM-1), particularly worrisome β -lactamases with the ability to inactivate most β -lactams, including carbapenems, and resistance to most marketed β -lactamase inhibitors.^{28,29} The structure of S-649266 (Fig. 2D) incorporates a catechol moiety in the 3-position substituent, which acts as a siderophore, and employs a 'Trojan Horse' strategy to increase cell penetration of the agent.³⁰ In this case, the catechol group acts as an iron chelator and enables the active transport of S-649266 into the bacterial cell by way of the cell's iron transport systems. S-649266 possesses the same dimethyl acetic acid substituted oxime group at the 7-position seen in ceftolozane and ceftazidime, which confers a high degree of β -lactamase resistance and activity against *P. aeruginosa*.²⁹ In fact, the compound has been reported to possess intrinsic resistance to the β -lactamases produced by many problematic strains of *Enterobacteriaceae*, including ESBL's, KPC, and NDM-1.³¹ Clinical trials currently underway are comparing S-649266 with imipenem/cilastatin for treatment of cUTI and are expected to be complete in September, 2016.³²

Next generation beta-lactamase inhibitors

Bacteria-produced β -lactamase enzymes hydrolytically cleave the cyclic amide bond of the β -lactam antibacterial agents before they are able to bind to and inhibit their target PBPs, representing a challenging bacterial drug resistance strategy. There are a large and diverse number of β -lactamase enzymes produced by bacteria (Table 1). The development of β -lactam antibacterials with intrinsic resistance to β -lactamases (*i.e.*, methicillin, higher generation cephalosporins, and the carbapenems) and mechanism-based inhibitors of the β -lactamase enzymes that can be co-administered with β -lactam antibacterials (Fig. 3), initially helped to overcome this drug resistance mechanism.⁸ Unfortunately, the recent emergence of inhibitor-resistant β -lactamases with activity against the previously intrinsically resistant β -lactams, has led to a critical lack of antibacterials with the capability of treating resistant infections. Particularly problematic β -lactamases that have seen recent emergence in Gram negative organisms include the extended-spectrum β -lactamases (ESBLs) such as TEM, SHV & CTX-M (class A), and OXA (class D), that have the ability to hydrolyze extended-spectrum cephalosporins with the oxime (oxymino) functional group.^{33–36} The AmpC-type β -lactamases (class C), seen in *Enterobacteriaceae* and *P. aeruginosa*, have the ability to hydrolyze broad and extended spectrum cephalosporins (cephamycins & oxime cephalosporins) and are not inhibited by classical β -lactamase inhibitors (*i.e.*, clavulanic acid).^{37,38} Lastly, the very recent emergence of carbapenemases, including KPC (class A), VIM, IMP, and NDM-1 (class B), is of great concern as these enzymes have the ability to hydrolyze

Table 1 Ambler classification of β -lactamases, substrate affinity, and selected examples. (Adapted with permission from Mandell's Principles & Practice of Infectious Disease, chapter 18, Table 18–2, p. 240)



Ambler class ^a	Enzyme activity	Substrates	Examples
A	Broad-spectrum (penicillinases)	Benzylpenicillins, aminopenicillins, carboxypenicillins, ureidopenicillins, narrow-spectrum cephalosporins	PC1 (<i>S. aureus</i>); TEM-1, SHV-1 (<i>E. coli</i> , <i>K. pneumoniae</i> & other G-ve bacteria)
A	Extended-spectrum (ESBL)	Substrates of broad-spectrum, plus: oxymino- β -lactams, aztreonam	TEM-derived, SHV-derived, CTX-M-derived (<i>Enterobacteriaceae</i>); PER-1, VEB-1, VEB-2, GES-1, GES-2, IBC-2 (<i>P. aeruginosa</i>)
A	Carbapenemases	Substrates of ESBLs, plus: cephamycins & carbapenems	KPC-1, KPC-2, KPC-3 (<i>K. pneumoniae</i>); NMC/IMI, SME family
B	Carbapenemases	Substrates of ESBLs, plus: cephamycins & carbapenems	NDM-1 (<i>Enterobacteriaceae</i>); IMP, VIM, GIM, SPM, SIM (<i>P. aeruginosa</i> , <i>Acinetobacter</i> spp.)
C	Cephalosporinases	Substrates of ESBLs, plus: cephamycins	AmpC-type (<i>Enterobacteriaceae</i> , <i>Acinetobacter</i> spp.)
D	Broad-spectrum (oxacillinases)	Aminopenicillins, ureidopenicillins, cloxacillin, methicillin, oxacillin, and some narrow-spectrum cephalosporins	OXA-family (<i>P. aeruginosa</i>)
D	Extended-spectrum (ESBL)	Substrates of broad-spectrum, plus: oxymino- β -lactams, aztreonam	OXA-derived (<i>P. aeruginosa</i>)
D	Carbapenemases	Substrates of ESBLs, plus: cephamycins & carbapenems	OXA-derived (<i>Acinetobacter</i> spp.)

^a Ambler class A, C, and D β -lactamases possess a catalytic active site serine; Ambler class B β -lactamases are Zn^{2+} metallo- β -lactamases.

the carbapenems that have been used as a last resort for treatment of organisms expressing ESBLs and AmpC.^{39–42} The recent, or near approvals of two new classes of β -lactamase inhibitors has provided hope of some reprieve from pan-drug resistant infections. The novel chemistry and mechanisms of these agents is discussed below.

Avibactam (formerly NLX104), a member of the diazabicyclo[3.2.1]octanone (DBO) series, is a novel β -lactamase inhibitor that does not possess the typical β -lactam ring seen in the previously marketed agents clavulanic acid, tazobactam, and sulbactam (Fig. 4A).^{43,44} Avibactam, in combination with the 3rd generation anti-pseudomonal cephalosporin, ceftazidime (Avycaz, Allergan, Inc.), was approved in the U.S. in 2015 for the treatment of cIAI (in combination with metronidazole), and cUTI caused by multidrug-resistant Gram negative bacteria. It is also in phase 2 clinical trials in Europe in combination with aztreonam for cIAIs and phase 3 trials in Europe for serious infections like complicated UTIs, acute pyelonephritis, and hospital acquired bacterial pneumonia.

Avibactam has an extended spectrum of activity against β -lactamase enzymes compared to the classical β -lactamase inhibitors, including activity against ESBLs, AmpC, and KPC β -lactamases.^{45,46} The combination of avibactam with ceftazidime extends the coverage of this agent to include activity against many resistant Gram-negative organisms, including ESBL-producing and carbapenem-resistant *Enterobacteriaceae*. The activity of the ceftazidime/avibactam combination against *P. aeruginosa* is modestly improved over ceftazidime alone, while *Acinetobacter* species remain mostly resistant to the combination drug.^{47,48} The activity of the combination product against anaerobic and Gram positive organisms is limited and similar to that of ceftazidime alone, which necessitates the combination with metronidazole for the treatment of cIAI.^{49,50} The structures of avibactam bound to class A (CTX-M), class C (AmpC), and class D (OXA) β -lactamases have been solved and provide key insights into

**Fig. 3** Marketed, mechanism-based β -lactamase inhibitors. A. Clavulanic acid. B. Tazobactam. C. Sulbactam.**Fig. 4** Next generation β -lactamase inhibitors recently approved or in clinical trials. A. Avibactam. B. Relebactam. C. Vaborbactam.

the drug's mechanism of β -lactamase inhibition.^{51–53} The classical β -lactamase inhibitors exert their activity by either transient or irreversible inhibition of the enzymes *via* the formation of an acyl-enzyme covalent bond associated with ring-opening. Hydrolysis of this bond can restore the active enzyme, but the β -lactamase inhibitor, once freed from the enzyme, is inactivated. Avibactam differs from the classical β -lactamase inhibitors by forming a carbamate linkage to the β -lactamase enzyme with opening of the diaza-bicyclo ring structure.⁵² The crystal structures indicate that the drug is more tightly bound within the β -lactamase active site compared with the classical inhibitors, with a large degree of rigidity along the carbamate linkage, and multiple hydrogen bonds in the active site.⁵⁴ Avibactam's displacement of a key water molecule in the active site that is presumed to be involved in the hydrolytic release of the classical β -lactamase inhibitors has been proposed to contribute to the drug's longer residence time in the active site compared with the classical inhibitors; *i.e.*, 50% of clavulanic acid is released from TEM-1 within 7 minutes, *versus* 7 days for avibactam.^{52,55} Lastly, upon release from the β -lactamase active site, avibactam has been shown to return to its original active form, unlike the classical β -lactamase inhibitors, which are inactive upon release.⁵⁶ Taken together, these factors contribute to the enhanced activity of avibactam over classical β -lactamase inhibitors against class A KPC β -lactamases, class C β -lactamases, and some class D β -lactamases.^{55,57}

Relebactam (formerly MK-7655, Merck) is another new β -lactamase inhibitor in the DBO class that is currently undergoing phase 3 clinical trials in Europe and the U.S. (Fig. 4B). The compound is being investigated for use in combination with imipenem–cilastatin (a carbapenem combined with a dehydropeptidase inhibitor that prevents degradation of imipenem in the kidney) for treatment of HAP, VAP, cIAI, and cUTI. Similar to avibactam, relebactam shows activity against class A and class C β -lactamases, including ESBLs.⁵⁸ When combined with imipenem–cilastatin, relebactam improved the activity of imipenem against carbapenemase-producing (KPC) *Enterobacteriaceae* and AmpC-producing *P. aeruginosa*.^{59,60} The combination is not effective against organisms expressing class B metallo- β -lactamases or class D carbapenemases and does not appear to add activity against *Acinetobacter baumannii*.^{61,62} Structurally similar to avibactam, relebactam has a piperidine ring added to the carbamoyl moiety at the 2-position. This compound was synthesized in an effort to test the effects of positively and negatively charged side chain substitutions to the carbamoyl group.⁶³ It is believed that a positively charged group at this position plays a key role in preventing the efflux of the drug from the bacterial cell, thereby facilitating synergy with imipenem. An X-ray crystal structure of relebactam bound in the active site of AmpC from *P. aeruginosa* (PDB entry 4NK3) shows the compound binds in a manner very similar to avibactam, with the piperidine substituent extending into an empty pocket of the active site and interacting primarily with water molecules (Fig. 5).⁶³




Fig. 5 Avibactam (cyan carbons) and relebactam (green carbons) shown bound into the active site of beta-lactamases, SHV-1 from *K. pneumoniae* [PDB ID 4ZAM] and AmpC from *P. aeruginosa* [PDB ID 4NK3], respectively.

In contrast to avibactam and relebactam, which possess a novel diaza-bicyclic-octanone scaffold, the novel β -lactamase inhibitor, vaborbactam (formerly RPX7009), possesses a unique, boronic acid-based cyclic scaffold (Fig. 4C). Vaborbactam is currently undergoing phase 3 clinical trials in Europe for use in combination with meropenem (Carbavance™, Rempex Pharmaceuticals) for treatment of cUTI, HAP, VAP, & bacteremia caused by carbapenem-resistant *Enterobacteriaceae*.^{64,65} Vaborbactam has been extensively studied in combination with the investigational carbapenem RPX2003 (biapenem).^{66,67} The combination showed excellent activity against class A carbapenemase-producing *Enterobacteriaceae*, including KPC-producing strains of *K. pneumoniae*. The boronic acid pharmacophore has been known for many years to have affinity for serine proteases,^{68,69} and boronic acid compounds have been explored as inhibitors of serine β -lactamases since the early 1970's.⁷⁰ A structure of a boronic acid-based compound covalently bound to a TEM β -lactamase was published in the early 90's,⁷¹ and led to over a decade of structure-guided design of boronic acid-based β -lactamase inhibitors.^{72–76} Vaborbactam differs from previously reported boronic acid inhibitors of β -lactamase by the rational design of a cyclic boronate ester, which was hypothesized to provide selectivity toward β -lactamase enzymes over other serine proteases, the latter showing preference for acyclic substrates due to a more sterically restricted active site.⁷⁷ Solved structures of vaborbactam bound to β -lactamases CTX-M-15 and AmpC show that the compound binds to the β -lactamase enzymes in a similar manner as avibactam and relebactam, with the exception that the catalytic serine residue forms a covalent bond with the boronic ester moiety (Fig. 6).⁷⁷

Next-generation oxazolidinone antibacterials

The oxazolidinone class of antibacterials, a fully synthetic class of agents, has been considered a major breakthrough in antimicrobial drug development.⁷⁸ The class is represented by the U.S. marketed agents linezolid (Zyvox®, Pfizer,




Fig. 6 RPX7009 bound to AmpC from *E. cloacae* [PDB ID 4XUX].

approved 2000) and tedizolid (Syvextro®, Merck, approved 2014), and several other promising agents currently in clinical development (Fig. 7). The history of the discovery and development of the oxazolidinone class as antibacterial agents have been previously reviewed,^{79–81} as well as early SAR studies.^{82–94} Oxazolidinones exert their antibacterial activity by binding to the bacterial 70S ribosome and inhibiting protein synthesis. The mechanism of this effect is unique from other classes of protein synthesis inhibitors in that they bind to the 50S ribosomal subunit, thereby preventing the binding

of aminoacyl-tRNA to the peptidyl transferase center at the A-site.^{95–97} The antibacterial spectrum of the oxazolidinone class is excellent with respect to most clinically relevant Gram positive pathogens, including methicillin-resistant *S. aureus* (MRSA), penicillin-resistant *S. pneumoniae* (PRSP), macrolide-resistant streptococci, and vancomycin-resistant enterococci (VRE).^{78,98} The class in general is clinically ineffective against most Gram negative pathogens, primarily due to an efflux mechanism of resistance.^{79,99–101} Linezolid is currently approved for use in the U.S. for treatment of nosocomial and community-acquired pneumonia (CAP), uncomplicated and complicated skin and skin structure infections (cSSSI), and infections caused by vancomycin-resistant enterococci. While a very promising agent for treatment of infections caused by resistant pathogens, linezolid is, unfortunately, not without its drawbacks. The drug is known to cause reversible thrombocytopenia and bone marrow suppression, neuropathies with prolonged use, and has been associated with adverse serotonergic effects due to its inhibition of monoamine oxidase (MAO).¹⁰² Further, the recent emergence of linezolid resistance has driven the investigation and advancement of next-generation oxazolidinones with improved antibacterial activities and decreased adverse effect profiles.^{103–105}

Tedizolid (formerly torezolid and TR-701), is a second-generation oxazolidinone that was approved in the U.S. in




Fig. 7 Oxazolidinones currently marketed or in clinical trials. A. Linezolid. B. Tedizolid. C. Radezolid. D. MRX-I. E. Cadezolid.

2014 for treatment of acute bacterial skin and skin structure infections (ABSSSI) caused by susceptible organisms (Fig. 7B).^{96,102,106–109} The drug is marketed as tedizolid phosphate, a prodrug formulation of tedizolid that is phosphorylated at the 5-position hydroxymethyl group.^{110–112} Structurally similar to linezolid, tedizolid possesses the core oxazolidinone ring (A-ring) as well as the fluorophenyl B-ring. As discussed in previous communications, the 5-R configuration of the A-ring and the *N*-aryl B-ring are minimally required for antibacterial activity.^{81,113} Fluorination of the B-ring improves antibacterial activity and this substituent is seen in all approved oxazolidinones as well as those in clinical development (Fig. 7). In tedizolid, a C-5 hydroxymethyl group replaces the acetamide moiety seen in linezolid. This group serves as an attachment point for the prodrug phosphate which improves water solubility and may decrease MAO inhibition. Additionally, the substitution of hydroxymethyl at this position decreases steric bulk which has been proposed to allow for retained activity against bacterial species with resistance mediated by active site methylation (*cfr* gene).^{113,114} Tedizolid also substitutes a bi-aryl system (C-ring pyridine and D-rings) for the morpholine C-ring seen in linezolid. This substitution has been suggested to be responsible for improved potency of tedizolid over linezolid, as the new ring system is able to form additional binding interactions in the 50S ribosomal subunit.^{115,116} With respect to the spectrum of activity, tedizolid is up to 8-fold more potent than linezolid against most Gram-positive staphylococci, streptococci, and enterococci, including MRSA and VRE species.^{117–121} It possesses moderately improved activity over linezolid against the Gram-negative *M. catarrhalis* and *H. influenzae*, an 8- to 16-fold improvement in activity against Gram-positive anaerobes, and 2-fold improvement in activity against Gram-negative anaerobes.^{117,120} Tedizolid also retains reasonable activity against linezolid-resistant Gram-positive cocci with *cfr*-mediated resistance.¹²² Additional favorable properties of tedizolid include a long half-life that allows once-daily administration (*versus* twice daily linezolid), low serotonergic effect due to decreased MAO interaction, and a significantly improved adverse event profile.^{96,102,106}

Radezolid (formerly RX-1741 and Rx-01), Melinta Therapeutics) (Fig. 7C) completed phase 2 clinical trials in the U.S. in 2009 for treatment of community-acquired pneumonia and uncomplicated skin and skin-structure infections.¹²³ The compound was rationally designed using experimental ribosome structures and computational models to have enhanced Gram-negative activity and improved oral bioavailability.^{124–127} Structurally, the drug replaces the D-ring of tedizolid with a (triazolyl-methyl)-aminoyl-methyl moiety attached *para* to a phenyl C-ring and possesses the same acetamide group at the 5-position of the oxazolidinone A-ring seen with linezolid.^{113,126–129} The antibacterial activity of radezolid has been evaluated in comparison to linezolid and the drug has been shown to have moderately improved activity against streptococci (*S. pneumoniae* and *S. pyogenes*), staphylococci (MSSA and MRSA), and enterococci, including

some activity against linezolid-resistant staphylococci, enterococci, and *S. pneumoniae*.¹²⁸ The safety profile and any advantages of radezolid compared to linezolid and tedizolid is not clear based on the current literature. To date, no phase 3 trials have been announced.

MRX-I (MicuRx Pharmaceuticals, Inc.) is another next-generation oxazolidinone currently in early phase 2 clinical trials in the U.S. for the treatment ABSSSI (Fig. 7D). This compound is reported to have been rationally designed to possess significantly decreased myelosuppression and inhibition of MAO resulting in an improved safety profile over linezolid.¹³⁰ MRX-I deviates from traditional SAR of the oxazolidinones by incorporating a novel 2,4,6-trifluorophenyl B-ring, a dihydropyridone C-ring, and a unique isoxazolyl-aminomethyl substitution at the A-ring 5-position. The trifluorophenyl B-ring is believed to be responsible for the improvement in the safety profile of MRX-I possibly due to a distortion of A-ring, B-ring planarity forced by the *ortho*-fluoro substituent that affects off-target binding to MAO and the mitochondrial ribosome.¹³⁰ With respect to the drug's spectrum of activity, MRX-I has similar antibacterial range to other oxazolidinones, including MRSA, PRSP, and VRE, and is comparable or slightly better than that of linezolid.¹³¹ The literature is not clear on the activity of MRX-I against linezolid-resistant bacterial strains.

The last agent of interest within the developing oxazolidinone class is cadazolid (ACT-179811, Actelion Pharmaceuticals, Ltd.).¹³² Structurally, cadazolid is almost a complete departure from molecules previously discussed. It is a bifunctional molecule containing the oxazolidinone core scaffold on one half tethered to a fluoroquinolone *via* a phenoxyether linkage (Fig. 7E).^{133–135} The common structural feature that serves as a bridge between the two molecules is a piperidine ring. This positioning of this ring is analogous to the morpholine seen in linezolid and the piperazine ring found in ciprofloxacin. Most reports seem to indicate that the primary mechanism of action for cadazolid is protein synthesis with very little effect on DNA synthesis as a possible secondary mechanism.¹³⁶ This would suggest that this agent acts more like an oxazolidinone with very little fluoroquinolone activity. It is being studied for the treatment of *C. difficile*-associated diarrhea (CDAD), and has progressed to phase 3 clinical trials in Europe and in the U.S. in comparison with vancomycin.^{137,138} It has very limited oral bioavailability due to its acidic and lipophilic nature, which means its activity is primarily localized to the gut.^{139,140} The drug has received QIDP and Fast Track development status from the U.S. FDA.

Next generation bacterial topoisomerase inhibitors

Currently, there are four independent drug classes that target the bacterial type II topoisomerases DNA Gyrase and topoisomerase IV. Chronologically, Novobiocin is the classic representative of the first class of such antibacterial agents. No longer commercially available, Novobiocin operates *via* the competitive inhibition of ATP on DNA Gyrase. Next, while

technically not the first class to exploit type II topoisomerases, the outstanding success of the quinolone class has almost single-handedly validated the target *via* the formation of DNA-quinolone-enzyme ternary complexes. Since their discovery in the 1960s, wide use of quinolones has resulted in high resistance levels. As such, rational improvements are continually being made to these indispensable agents, some of which are discussed below. Thirdly, a novel class of bacterial type II topoisomerase inhibitor has arisen that occupies a topoisomerase binding site distinctly separate from the quinolones and, therefore, possesses a unique MOI. These agents, known as novel bacterial topoisomerase inhibitors (NBTIs), consequently lack cross-resistance among pathogens possessing quinolone resistance. Lastly, the newer spiropyrimidinetrione class has recently arisen with yet another unique mechanism of topoisomerase II inhibition. Though discovered *via* whole-cell activity screens, spiropyrimidinetriones have since been rationally optimized.¹⁴¹ As one would expect, the unique MOIs of these newer classes imply accordingly unique structures, scaffolds, and SARs.

Zoliflodacin (previously known as ETX0914/AZD0914, Entasis Therapeutics, Inc.) is an oral antibacterial agent with QIDP fast-track status currently in phase 2 clinical trials in the U.S. for the treatment of uncomplicated gonorrhea. It is a spiropyrimidinetrione, a novel class of topoisomerase II inhibitors that uniquely inhibits re-ligation and consequently

causes a bactericidal buildup of DNA double-strand cleavages.^{142,143} It shows no cross-resistance with other type II topoisomerase inhibitor classes and exhibits an antibacterial spectrum of activity (SOA) that includes Gram-positive organisms; fastidious Gram-negative organisms such as *Neisseria gonorrhoeae*, *Haemophilus influenzae*, and *Moraxella catarrhalis*; as well as anaerobes.^{142–144} While much of the SAR was previously optimized or intolerable to manipulation, the benzisoxazole 3-position on the spiropyrimidinetrione scaffold proved to be modifiable. Thorough SAR optimization at this position resulted in the minimization of toxicity issues seen with fluoroquinolones (Fig. 8A).¹⁴²

Gepotidacin (also known as GSK2140944, GlaxoSmithKline plc) is a novel NBTI currently in phase 2 clinical trials in Europe for uncomplicated urogenital gonorrhea and the U.S. for Gram-positive ABSSSI, respiratory tract infections, and uncomplicated gonorrhea. It is a triazaacenaphthylene that binds to both *gyrA* and *parC* subunits at sites distinctly different from the fluoroquinolones and aminocoumarin-binding sites.^{145,146} Early NBTIs arose from whole cell antibacterial screens and have since been structurally characterized.^{147,148} Structural motifs include a bicyclic aromatic heterocycle connected to another aromatic heterocycle by a variable linker that must contain a basic nitrogen at position-7 in order to form a requisite salt bridge with Asp83 as seen *via* X-ray crystallographic structural data (Fig. 8B).^{146,147} NBTIs commonly exhibit high potencies and rather broad-




Fig. 8 Next generation bacterial topoisomerase inhibitors. A. Zoliflodacin. B. Gepotidacin. C. Finafloxacin. D. Nemonoxacin. E. Delafloxacin. F. Zabofloxacin. G. Avarofloxacin.

spectrums of activity, but are generally plagued by unacceptable hERG activity that appears to be determined by overall compound polarity.¹⁴⁶ Gepotidacin, however, has shown relatively low hERG activity with an IC₅₀ of 1.4 mM.¹⁴⁶

Finafloxacin (MerLion Pharmaceuticals Pte Ltd., Singapore; licensed as Xtoro® under Alcon Pharmaceuticals in the U.S.) is an 8-cyano fluoroquinolone with QIDP status that, while already approved in the U.S. in the form of an otic suspension for the treatment of acute otitis externa, is currently in phase 2 clinical trials in the U.S. for the treatment of ventilator-associated bacterial pneumonia (VABP) due to pseudomonas, acute bacterial skin and skin structure infections (ABSSSI), and complicated intra-abdominal infections. Unlike other fluoroquinolones, which exhibit reduced activity under acidic conditions, finafloxacin shows improved activity under slightly acidic conditions (pH 5.0–6.0) due to a unique 7-substituent chiral base component that confers a distinctive base capacity (Fig. 8C).^{149,150} It is presumed that the stereo-configuration of the 7a hydrogen on the base component is responsible for this characteristic. Finafloxacin's ability to act under harsh acidic environments grants it a number of potential applications, such as *Helicobacter pylori* infections, urinary tract infections (UTIs), and more.

Nemonoxacin (TaiGen Biotechnology Co., Ltd.) is a non-fluorinated C-8-methoxy quinolone with QIDP status that has entered phase 3 clinical trials for the treatment of community acquired pneumonia (CAP), ABSSSI, and diabetic foot infection. The C-8-methoxy group grants nemonoxacin activity against both DNA Gyrase and topoisomerase IV, thereby decreasing resistant mutant selection, while the removal of the C-6-fluorine is presumed to decrease the incidence of toxic side effects (Fig. 8D).^{151,152} As compared to standard fluoroquinolones, it has shown increased activity against certain Gram-positive pathogens, including MSSA, MRSA, and multi-drug resistant *S. pneumonia*, but generally shows less activity against Gram-negative pathogens, including similar or inferior results as compared to standard fluoroquinolones.^{152,153} Nemonoxacin has also shown varying degrees of activity against different quinolone-resistant pathogens.¹⁵⁴

Delafloxacin (Baxdela™, Melinta Therapeutics) is a fluoroquinolone with QIDP status in phase 3 clinical trials for the treatment of ABSSSI and at the time of this review, is soon set to begin phase 3 trials for the treatment of CAP. Unlike other fluoroquinolones, delafloxacin exhibits a similar affinity for both DNA Gyrase and topoisomerase IV in both Gram-positive and Gram-negative pathogens and, therefore, is expected to be less selective for resistance mutants.^{155,156} Delafloxacin has not been found to be a more potent topoisomerase inhibitor than other fluoroquinolones, but has been found to have greater potency, particularly against Gram-positive bacteria.¹⁵⁶ Structural characteristics implicated for this attribute include a bulky heteroaromatic substituent (6-amino-3,5-difluoropyridine) at the 1 position, a C-8-chlorine that grants weak polarity, and the lack of a basic group at the 7 position (3-hydroxyazetidine) (Fig. 8E).¹⁵⁶ This last characteristic grants delafloxacin properties particularly

unique among quinolones. Lacking a basic group available for protonation at the 7 position results in an anionic charge at neutral pH values (whereas most quinolone antibacterials form zwitterions at a physiological pH) and a mainly neutral charge at a slightly acidic pH.¹⁵⁶ As such, delafloxacin has shown improved activity under slightly acidic conditions, which as mentioned above, is a rare characteristic for quinolones, and results in improved MICs against different bacteria as compared to moxifloxacin and ciprofloxacin.^{157–159}

While there is less rational design information readily available at the time of this review, two more quinolones are being developed and in late-phase trials. Zabofloxacin (Dong Wha Pharmaceutical Co., Ltd.) is a fluoroquinolone that has completed phase 3 trials for the treatment of acute bacterial exacerbations of chronic obstructive pulmonary disease (COPD) (Fig. 8F). Avarofloxacin (previously known as JNJ-Q2 and JNJ-32729463, Janssen Pharmaceutica, licensed to Furix Pharmaceuticals) is a wide-spectrum fluoroquinolone with QIDP status that has completed phase 2 trials for the treatment of complicated skin and skin structure infections (CSSSIs) (Fig. 8G).

Next generation tetracyclines

Since their discovery in the 1940s, tetracyclines have been widely used to treat both Gram-positive and Gram-negative pathogens. SAR studies of tetracycline analogues and the co-crystallization of tetracycline with the 30S subunit revealed significant interactions between several functional groups in the drug and nucleotides in the ribosomal structure (Fig. 9).¹⁶⁰ The same studies showed a lack of significant interactions between ring D and ribosomal nucleotides. The lack of significant binding at the C-7 through C-9 positions spurred the search for ring D-modified analogs, with the resulting development of tigecycline. Tigecycline (Tygacil®; Wyeth Pharmaceuticals Inc., a subsidiary of Pfizer Inc., initial U.S. approval in 2005) is the first semi-synthetic representative of a class of antimicrobials called glycylcyclines, which are 9-*t*-butylglycylamido derivatives of minocycline, a second generation tetracycline (Fig. 10A).¹⁶¹ The design and synthesis of glycylcyclines were intended to overcome the major problems of resistance associated with the tetracyclines, namely ribosomal protection and efflux-pump-mediated resistance.^{162,163} Tigecycline consequently demonstrates good *in vitro* and *in vivo* activity against a wide range of mutant bacterial strains enabled with these resistance mechanisms, as well as potent activity against MRSA, penicillin-resistant *streptococci* (PRS), and VRE.^{160,162,164} SAR studies determined that the *t*-butylglycylamido moiety of tigecycline confers the highest activity, and replacement of glycine with other amino acids does not further enhance it.¹⁶⁵ The 7-deaminated analog of tigecycline showed comparable antimicrobial activity.¹⁶⁶ The binding mode of tigecycline to the 70S ribosome has been elucidated as well, showing similar interactions as observed with tetracycline, but with some significant differences.¹⁶⁷ In




Fig. 9 Tetracycline (cyan carbons) shown binding to the bacterial ribosome, 30S subunit [PDB ID 1HNW]. Hydrogen bonds are shown as dashed, yellow lines.




Fig. 10 Next generation tetracyclines & glycylcyclines marketed or in clinical trials. A. Tigecycline. B. Omadacycline. C. Eravacycline.

particular, although both the 7-dimethylamino and the 9-*t*-butylglycylamido moieties of tigecycline do not seem to directly interact with any ribosomal nucleotide, they may enhance binding by increasing electron density of ring D, thereby enhancing the π -stacking interaction with C1054. The rigidity of the 9-*t*-butylglycylamido moiety also seems to induce conformational changes that allow stacking between C1054 and U1196, similar to the ribosomal conformation when tRNA is bound to the A site. The success and utility of tigecycline, coupled with climbing resistance rates against other major antibacterial classes, have recently prompted the development of several more novel tetracycline analogues.

Omadacycline (Paratek Pharmaceuticals) is a novel tetracycline derivative with QIDP status in phase 3 trials for the treatment of ABSSSIs and CAP. Specifically, omadacycline is a C-9-substituted semi-synthetic minocycline derivative known as an aminomethylcycline (AMC) (Fig. 10B).^{168,169} It is a broad-spectrum antibacterial that shows strong activity against various tetracycline-resistant Gram-positive and Gram-negative organisms.^{168,169} It has shown *in vitro* activity against additional organisms as well, including anaerobes, atypical pathogens, and organisms resistant to other classes of antibacterials, such as methicillin, vancomycin and ciprofloxacin.¹⁶⁸ During the development of omadacycline,

diversity sets based around a 9-aminomethylminocycline intermediate were analyzed to find and explore hits against tetracycline-resistant Gram-positive organisms.¹⁶⁹ Initially increasing alkyl group size at this C-9 position was found to substantially increase activity against these organisms, while the side chain seen at this position on omadacycline was found to be optimal for activity as further increased methyl branching and different alkyl chain length were both found to decrease activity.¹⁶⁹ Omadacycline is being formulated in both oral tablet form and intravenous form.¹⁶⁸

Eravacycline (Tetraphase Pharmaceuticals) is another next-generation tetracycline, known as a fluorocycline. It has QIDP status and is in phase 3 trials for the treatment of cUTIs and cIAIs. While semi-synthetic development of tetracyclines has previously been hindered by a limited ability to manipulate C-7 and C-9 substituents, total synthesis approaches have recently been developed that allow for improved diversity at these positions (as well as C-8), which has allowed for the development of new 7,9-disubstituted tetracycline analogues, including eravacycline.¹⁷⁰ As electronegativity at the C-7 substituent has been shown to effect the antibacterial potency of tetracyclines, a fluoro substituent at this position is found on eravacycline (Fig. 10C).^{170,171} Furthermore, SAR studies showed a small alkylamine substituent on the C-9 side chain to be the optimum amino group for antibacterial activity.¹⁷⁰ Eravacycline has shown broad-spectrum *in vitro* activity against aerobic and anaerobic Gram-positive and Gram-negative pathogens, as well as MDR pathogens including ESBL- and carbapenemase-producing pathogens.¹⁷² It also maintains activity against tetracycline efflux pumps and ribosomal protection.¹⁷³ Additionally, while results are still being analyzed, it is reported that eravacycline did not achieve its primary endpoint of statistical non-inferiority compared to levofloxacin for cUTI in the phase 3 trial IGNITE2.¹⁷⁴

Next generation macrolides

Macrolide antibiotics are natural product-based antibacterials that are composed of a large, macrocyclic lactone core, to which a variety of sugar substituents may be attached. Erythromycin, which was discovered in soil samples collected in the Philippine islands, entered the U.S. market in 1952 and was followed by a variety of other macrolides over the following decades, notably clarithromycin (Biaxin, approved 1995) and azithromycin (Zithromax, approved 1996). Macrolide antibiotics exert their antibacterial effect by binding to and inhibiting the action of the bacterial 70S ribosome, specifically by binding to the 23S rRNA component of the 50S subunit and subsequently inhibiting cellular protein synthesis. The macrolide antibiotics are notable for their Gram-positive activity and are typically prescribed to treat respiratory tract infections caused by Gram-positive organisms (*Streptococci*, *Staphylococci*, and *Enterococci*) as well as certain “atypical” organism, including *Legionella pneumophila*, *Mycoplasma* spp., *Chlamydia* spp. and some *Rickettsia*. Disadvantages of the macrolide antibiotics include limited Gram-

negative activity, acid instability limiting the oral effectiveness of some compounds in the class, and interaction with human drug metabolizing enzymes in the cytochrome P-450 system that can result in significant drug interactions.¹⁷⁵ Further, emerging resistance, typically due to alterations of the macrolide binding site on the ribosome, has limited the clinical utility of older compounds in the class.

Telithromycin (Ketek, Sanofi-Aventis) was the first ‘next-generation’ macrolide to be approved for use in the U.S. in 2004 (Fig. 11A). The compound, the first in the ‘ketolide’ sub-class, is a semi-synthetic erythromycin derivative that substitutes a keto functionality at the 3-position for an L-cladinose. Additionally, a novel carbamate ring has been attached to the central lactone system at the 11- and 12-positions with a linked aryl-alkyl moiety.¹⁷⁶ The 3-keto functionality is believed to be responsible for several desirable properties of the ketolides, including a higher degree of acid stability (aided by the 6-position methyl group), a lack of induction of macrolide resistance (MLS_B phenotype), and the ability to overcome resistance caused by methylation of the 23S subunit (a component of the 50S subunit).^{176,177} Further, the carbamate substituent plays a role in the expanded activity of telithromycin compared to erythromycin, and has been suggested to mediate the drug's resistance to bacterial efflux and play a role in improved pharmacokinetic properties.^{100,101,176,178} Classical macrolides interact with domain V of the 23S rRNA subunit. The increased binding affinity of telithromycin has been shown to be due to an additional interaction between the carbamate functionality of telithromycin and residues in domain II of the 23S subunit.¹⁷⁹ This increased binding affinity presumably allows the ketolide antibiotics to overcome resistance due to active site methylation and confers activity of telithromycin against erythromycin-resistant Gram-positive organisms. The drug shows excellent activity against most Gram-positive aerobic organisms, including macrolide-resistant strains of *S. pneumoniae*, excellent activity against atypical pathogens including *C. pneumoniae*, *L. pneumophila*, and *M. pneumoniae*, and good activity against some Gram-negative aerobes including *M. catarrhalis* and *H. influenzae*.^{177,180–182} Although initially showing great promise and receiving U.S. approval for use in treatment of community-acquired pneumonia, acute bacterial sinusitis, and bacterial exacerbations of chronic bronchitis, telithromycin was subsequently discovered to cause irreversible hepatotoxicity in some patients and the sinusitis and bronchitis indications were removed by the FDA in 2007 and a black box warning added to the package literature.¹⁸³ Clinical trial safety and data integrity concerns with this drug have been raised and have somewhat overshadowed the promise of the ketolide class.^{184,185}

Two new ketolide antibiotics that are currently in or have completed U.S. clinical trials include cethromycin (Fig. 11B) (Restanza, ABT-773, Advanced Life Sciences, Inc.) and solithromycin (Fig. 11C) (CEM-101, Cemptra, Inc.). Cethromycin completed phase 3 clinical trials in 2007 for treatment of community-acquired pneumonia. However,




Fig. 11 Advanced generation macrolides. A. Telithromycin. B. Cethromycin. C. Solithromycin.

while it was found to be safe, the drug was denied U.S. approval by the FDA in 2009 for reasons related to clinical trial design.¹⁸⁶ Structurally, cethromycin possesses the 3-position keto group characteristic of the ketolide class as well as the 11, 12 carbamate moiety. Similar to telithromycin, cethromycin possesses an aryl-alkyl side chain, but in this case it is a quinolylallyl side chain attached to the C6 position *versus* the carbamate nitrogen in telithromycin. These structural features impart pharmacology similar to telithromycin, with two points of interaction with the 23S rRNA subunit, at domains II and V, affording the drug a spectrum of activity similar to telithromycin including activity against erythromycin resistant organisms and some resistance to efflux.^{187,188} Solithromycin (CEM-101, Cempira, Inc.) is a novel fluoroketolide in phase 2 clinical trials in Europe for COPD and has recently completed phase 3 clinical trials in the U.S. comparing it to moxifloxacin for the treatment of community-acquired pneumonia.^{189,190} Solithromycin is structurally similar to telithromycin, with a 3-keto moiety, an 11,12 carbamate with an aryl-alkyl substitution (in this case a butyl-[1,2,3]-triazolyl-aminophenyl), and a unique fluorine substitution at the 2-position of the lactone (Fig. 11C). With respect to its spectrum of activity, solithromycin shows a similar profile to telithromycin, with enhanced activity against some telithromycin intermediate and resistant organisms.¹⁹¹ The enhanced activity has been proposed to be due to improved binding of solithromycin to the bacterial ribosome over telithromycin, including Erm-methylated ribosomes.¹⁹² The X-ray crystal structure of solithromycin bound to the *E. coli* ribosome provides insight into possible reasons for this improved binding.¹⁹² First, the aryl-alkyl side chain forms interactions with rRNA base pairs in domain II of the 23S subunit, including a possible hydrogen bond mediated by the amino group. Second, weak electrostatic interactions of the 2-fluorine with an rRNA base near domain V may enhance the activity of solithromycin over telithromycin. Lastly,

as with other ketolides, the substitution of the keto group at the 3-position for the cladinose sugar seen in earlier macrolide generations contributes to the binding affinity of the drug for Erm-methylated bacterial ribosomes.

Next generation aminoglycosides

Plazomicin (ACHN-490, Achaogen, Inc.) is a next generation, semi-synthetic aminoglycoside (neoglycoside) with reported activity against a broad range of Gram-positive and Gram-negative organisms, including carbapenem-resistant *Enterobacteriaceae*, when used in combination therapy with other antibacterial agents.¹⁹³ Derived by synthetic modification of sisomicin, a hydroxy-aminobutyric acid substituent at position 1 and a hydroxyethyl substituent at position 6' are reported to protect the compound against aminoglycoside-modifying enzymes in clinically relevant organisms (Fig. 12).¹⁹⁴ Plazomicin is not active against organisms with aminoglycoside resistance due to the expression of ribosomal methyltransferases. This agent is currently undergoing phase 3 clinical trials in Europe for blood stream infections and nosocomial pneumonia due to carbapenem-resistant *Enterobacteriaceae* as well as cUTIs and acute pyelonephritis. It is also in phase 3 trials in the U.S. for treatment of complicated urinary tract infections, hospital- and ventilator-associated bacterial pneumonia, and complicated intra-abdominal infection.

Novel bacterial folate synthesis inhibitors

Iclaprim (Motif Bio plc) is a next generation dihydrofolate reductase (DHFR) inhibitor with QIDP status in phase 3 clinical trials for ABSSSI and cSSSI. It is a novel 2,4-diaminopyrimidine trimethoprim analogue that was rationally designed and synthesized by F. Hoffman-La Roche Ltd. using structural data and molecular modeling (Fig. 13).¹⁹⁵ Studies show a single point mutation on DHFR to be




Fig. 12 Plazomicin.

responsible for trimethoprim resistance, illustrated by an F98Y mutation in *S. aureus* strain B71 that causes the loss of a single H-bond between trimethoprim's 4-amino group and the enzyme, resulting in a 64-fold increase in MIC.¹⁹⁶ It is posited that increased hydrophobic interactions between DHFR and iclaprim increase affinity, thereby overcoming trimethoprim resistance. Structural data shows that within the hydrophobic channel of DHFR, positioning of the *p*-aminobenzoic acid moiety of folate is similar to that of the chromene moiety of iclaprim.¹⁹⁷ It has potent Gram-positive bactericidal activity, including activity against MSSA, MRSA, and trimethoprim-resistant F98Y mutant strains.¹⁹⁷ Activity against Gram-negative and atypical organisms is reported to

be similar to that of trimethoprim.¹⁹⁸ The drug has reported synergy with the sulfonamide inhibitors of dihydropteroate synthase (DHPS), another key enzyme in the bacterial folate pathway.^{199,200} In clinical trials, iclaprim has been compared with linezolid as monotherapy for cSSSI (ASSIST-1 and ASSIST-2 trials) and further phase 3 trials are underway comparing the drug, again as monotherapy, with vancomycin for ABSSSI (REVIVE-1 and REVIVE-2 trials).³² With respect to selectivity over mammalian DHFR, iclaprim has been found to inhibit bacterial DHFR at sub-micromolar concentrations and lack inhibition of human DHFR at more than 5 orders of magnitude higher concentrations, showcasing favorable selectivity.¹⁹⁵

Novel inhibitors of the bacterial FAS-2 pathway

The type II bacterial fatty acid synthesis pathway, FAS II, provides fatty acid precursors for membrane phospholipids that are essential to bacterial cells. The pathway is distinct from the mammalian FAS I pathway which is composed of a single, multifunctional synthase, thus the FAS II enzymes represent novel and selective antibacterial targets that remain relatively unexploited. The structure and function of the enzymes in the bacterial FAS II pathway have been previously reviewed.^{201–204} As a metabolic pathway there is some concern about the ability of bacteria to bypass FAS II inhibition using exogenous fatty acids and this has been the subject of intense debate.^{205–207} Recent work, however, has demonstrated that certain bacteria, including *S. aureus*, remain susceptible to FAS II inhibition even in the presence of exogenous fatty acids due to differences in their regulation of genetic expression and feedback regulatory systems.^{208,209} As a rate-limiting enzyme in the FAS-II pathway, FabI, enoyl-[acyl-carrier protein (ACP)] reductase, represents a particularly attractive drug target and a number of inhibitors of this enzyme have been characterized.^{210–221} Triclosan, a stereotypical FabI inhibitor with a diphenyl ether scaffold, has been marketed for a number of years and is used in a variety of over-the-counter household products as a sterilizing agent, however it has low utility for systemic use because of poor bioavailability.^{222,223} Two promising FabI inhibitors, Debio 1452 and CG400549, with activity against *S. aureus*, have progressed sufficiently in clinical trials to merit discussion here.

Debio 1452 (previously AFN-1252, Debiopharm Group) is a FabI inhibitor with specific activity against staphylococci, including MRSA.^{219,224} The drug has little to no activity against non-staphylococci Gram-positive organisms, including streptococci, due to the ability of these organisms to bypass the FAS II pathway with exogenous fatty acids or the absence of the FabI target enzyme in some species such as the streptococci.²⁰⁸ Further, Debio 1452 has negligible Gram-negative activity, making the drug quite selective for *S. aureus* and coagulase-negative staphylococci. The drug has recently completed U.S. phase 2 clinical trials for ABSSSI with favorable outcomes.²²⁵ A produg formulation, Debio 1450, allowing




Fig. 13 Bacterial DHFR inhibitors. A. Trimethoprim. B. Iclaprim.




Fig. 14 Inhibitors of the bacterial FAS2 pathway. A. Debio 1452. Debio 1450 is a reported prodrug of Debio 1452, with undisclosed structure. B. CG-400549.

for oral and IV formulations is also currently in phase 2 clinical trials for ABSSSI.²²⁶ Structurally, Debio 1452 consists of a 3-methyl-benzofuran and an oxo-tetrahydro-naphthyridine linked by an *N*-methyl-propenamide (Fig. 14A).²²⁷ The compound was initially discovered using an iterative, structure-guided strategy, and has an interesting mechanism of inhibition involving the drug binding to a binary complex of enzyme and oxidized cofactor, NADP⁺.^{219,227,228}

CG400549 (Crystal Genomics, Inc.) completed a phase 2 clinical trial in 2012 for treatment of cABSSSI caused by MRSA.³² Similar to Debio 1452, CG400549 has selective anti-staphylococcal activity, including activity against MRSA.^{229–231} The company announced the completion of the phase 2 trial and confirmed *in vivo* efficacy with minimal adverse events, however no plans for subsequent clinical trials have been reported.²³² Structurally, the compound is a 2-pyridone analog (Fig. 14B), having been rationally designed to improve on the poor pharmacokinetics of the diphenyl ether compounds represented by triclosan, specifically by replacing a metabolically unstable phenol group of triclosan with the pyridone ring system and the ether functionality of triclosan with a methylene group.^{215,232} Recent X-ray structures have revealed the binding interactions of CG400549 in *E. coli* FabI and *S. aureus* FabI.²³² The keto group of CG400549's pyridone ring, similar to the amide keto in Debio 1452, forms electrostatic interactions with both the protein (a key tyrosine residue) and the NADPH cofactor. Unlike Debio 1452, CG400549 is believed to bind to the protein–NADPH (reduced cofactor) binary complex, making the drug uncompetitive with respect to NADPH (*vs.* NADP⁺ for Debio 1452) and competitive with respect to the enoyl substrate. This structural information has guided recent efforts to expand the activity of CG400549 by iterative, structure-guided design strategies, to include Gram-negative pathogens and mycobacteria.²³²

Conclusions

The number of novel antibacterials recently approved or under late-stage development is highly encouraging, as is recent U.S. and European legislation facilitating the rapid development and approval of new antibacterials. At the time of this review, there are almost 60 antimicrobial drugs that have gained QIDP status in the U.S., 6 of which have already attained U.S. approval, and 37 that are in late-phase clinical

trials.²³³ These results stand as evidence for the initial success of these new initiatives and their ability to drive the advancement of antibacterial development. A thorough analysis of the antibacterial pipeline, however, showcases the relatively low number of compounds under development with original targets, thereby highlighting the urgent need for the characterization and validation of such. Furthermore, it has been argued that ample attention should be given to the future development of selective antibacterial targets, giving way to what some predict will be the next phase of antibacterial discovery and development, the “narrow spectrum-era”.¹ It is reasonable to believe that antibacterial agents geared toward such selective targets would have substantial benefits—such as the possibility of developing effective microbiome-sparing and resistance-mitigating therapeutic strategies. Characterizing and validating novel narrow-spectrum targets, of course, would require an understandably significant amount of resources, innovation, and effort—including fully rational drug design based around heretofore unconventional targets, the efficient use of modern techniques like diversity-oriented synthesis, as well as a certain degree of *anticipated* success.

Current techniques and methods being implemented in the rational design and optimization of antibacterial agents forecast a possible change in the state of affairs regarding infectious disease. As described above, recent advances in various techniques have helped gain ground in a struggle against bacterial infections where prolonged trends have demonstrated mostly the opposite. Promising examples include specific chiral configurations of new quinolones that grant improved activity under acidic conditions, fully-synthetic techniques used in the next-generation tetracyclines that allow for novel side-chains that grant activity against resistant pathogens, and semi-synthetic techniques used in the next-generation macrolides that allow for the modification of functional groups that yields subsequent binding improvements and, again, activity against resistant pathogens. Upon review, the recent advances in the rational design and optimization of antibacterial agents are, for the time being, highly encouraging.

Competing interests

The authors declare no competing interests regarding the content of this manuscript.

Abbreviations

ABSSSI	Acute bacterial skin and skin structure infection
AMC	Aminomethylcycline
CAP	Community acquired pneumonia
cIAI	Complicated intra-abdominal infection
COPD	Chronic obstructive pulmonary disease
cSSSI	Complicated skin and skin structure infection
DHFR	Dihydrofolate reductase
GAIN act	Generating antibiotics incentives now act
HTS	High-throughput screening
IMI	Innovative medicines initiative
MOI	Mode of inhibition
MRSA	Methicillin-resistant <i>Staphylococcus aureus</i>
MSSA	Methicillin-sensitive <i>Staphylococcus aureus</i>
NBTIs	Novel bacterial topoisomerase inhibitors
ND4BB	New Drugs 4 Bad Bugs
QIDP	Qualified infectious disease product
SOA	Spectrum of activity
VABP	Ventilator-associated bacterial pneumonia

Acknowledgements

KH gratefully acknowledges ITHS Rising Stars Career Development Program at University of Washington, which is supported by grants UL1TR000423, KL2TR000421, and TL1TR000422 from the NIH National Center for Advancing Translational Sciences through the Clinical and Translational Science Awards Program (CTSA). KH also acknowledges the support of the Mountain West IDeA Clinical and Translational Research – Infrastructure Network (CTR-IN), which is supported by a grant from the National Institute of General Medical Sciences (5 U54 GM104944). Support from the Idaho State University faculty development programs (Office of Research and College of Pharmacy) is also gratefully acknowledged.

References

- 1 E. D. Brown and G. D. Wright, *Nature*, 2016, **529**, 336–343.
- 2 Review on Antimicrobial Resistance: Antimicrobial Resistance: Tackling a Crisis for the Health and Wealth of Nations, 2014.
- 3 Centers_for_Disease_Control_and_Prevention, Antibiotic resistance threats in the United States, 2013, Accessed June 17, 2015.
- 4 M. A. Cooper and D. Shlaes, *Nature*, 2011, **472**, 32.
- 5 D. J. Payne, M. N. Gwynn, D. J. Holmes and D. L. Pompliano, *Nat. Rev. Drug Discovery*, 2007, **6**, 29–40.
- 6 K. Lewis, *Nat. Rev. Drug Discovery*, 2013, **12**, 371–387.
- 7 L. L. Ling, T. Schneider, A. J. Peoples, A. L. Spoering, I. Engels, B. P. Conlon, A. Mueller, T. F. Schaberle, D. E. Hughes, S. Epstein, M. Jones, L. Lazarides, V. A. Steadman, D. R. Cohen, C. R. Felix, K. A. Fetterman, W. P. Millett, A. G. Nitti, A. M. Zullo, C. Chen and K. Lewis, *Nature*, 2015, **517**, 455–459.
- 8 S. A. Testero, J. F. Fisher and S. Mobashery, in *Burger's Medicinal Chemistry and Drug Discovery*, ed. D. J. Abraham and D. P. Rotella, John Wiley and Sons, Inc., Hoboken, NJ, 7th edn, 2010, vol. 7, pp. 257–402.
- 9 T. Ishikawa, N. Matsunaga, H. Tawada, N. Kuroda, Y. Nakayama, Y. Ishibashi, M. Tomimoto, Y. Ikeda, Y. Tagawa, Y. Iizawa, K. Okonogi, S. Hashiguchi and A. Miyake, *Bioorg. Med. Chem.*, 2003, **11**, 2427–2437.
- 10 R. K. Flamm, H. S. Sader, D. J. Farrell and R. N. Jones, *Antimicrob. Agents Chemother.*, 2012, **56**, 2933–2940.
- 11 H. S. Sader, D. J. Farrell, R. E. Mendes, R. K. Flamm, M. Castanheira and R. N. Jones, *Diagn. Microbiol. Infect. Dis.*, 2015, **82**, 78–84.
- 12 E. J. Goldstein, D. M. Citron, C. V. Merriam and K. L. Tyrrell, *Diagn. Microbiol. Infect. Dis.*, 2013, **76**, 347–351.
- 13 G. G. Zhanel, G. Snizek, F. Schweizer, S. Zelenitsky, P. R. Lagace-Wiens, E. Rubinstein, A. S. Gin, D. J. Hoban and J. A. Karlowsky, *Drugs*, 2009, **69**, 809–831.
- 14 J. L. Del Pozo and R. Patel, *Drugs of today*, 2008, **44**, 801–825.
- 15 H. Canada, *Zeftara (ceftobiprole medocartil) for Injection - Discontinuation of Sale - Notice to Hospitals*, <http://www.healthycanadians.gc.ca/recall-alert-rappel-avis/hc-sc/2010/14632a-eng.php>, Accessed June 14, 2016.
- 16 T. R. Fritsche, H. S. Sader and R. N. Jones, *Diagn. Microbiol. Infect. Dis.*, 2008, **61**, 86–95.
- 17 C. Bustos and J. L. Del Pozo, *Infect Drug Resist*, 2010, **3**, 5–14.
- 18 E. J. Goldstein, D. M. Citron, C. V. Merriam, Y. A. Warren, K. L. Tyrrell and H. T. Fernandez, *Antimicrob. Agents Chemother.*, 2006, **50**, 3959–3962.
- 19 N. C. Issa, M. S. Rouse, K. E. Piper, W. R. Wilson, J. M. Steckelberg and R. Patel, *Diagn. Microbiol. Infect. Dis.*, 2004, **48**, 73–75.
- 20 A. L. Lovering, M. C. Gretes, S. S. Safadi, F. Danel, L. de Castro, M. G. Page and N. C. Strynadka, *J. Biol. Chem.*, 2012, **287**, 32096–32102.
- 21 H. S. Sader, P. R. Rhomberg, D. J. Farrell and R. N. Jones, *Antimicrob. Agents Chemother.*, 2011, **55**, 2390–2394.
- 22 D. M. Livermore, S. Mushtaq and Y. Ge, *The Journal of antimicrobial chemotherapy*, 2010, **65**, 1972–1974.
- 23 E. S. Armstrong, D. J. Farrell, M. Palchak and J. N. Steenbergen, *Antimicrob. Agents Chemother.*, 2015, **60**, 666–668.
- 24 A. Toda, H. Ohki, T. Yamanaka, K. Murano, S. Okuda, K. Kawabata, K. Hatano, K. Matsuda, K. Misumi, K. Itoh, K. Satoh and S. Inoue, *Bioorg. Med. Chem. Lett.*, 2008, **18**, 4849–4852.
- 25 K. Murano, T. Yamanaka, A. Toda, H. Ohki, S. Okuda, K. Kawabata, K. Hatano, S. Takeda, H. Akamatsu, K. Itoh, K. Misumi, S. Inoue and T. Takagi, *Bioorg. Med. Chem.*, 2008, **16**, 2261–2275.
- 26 S. Takeda, Y. Ishii, K. Hatano, K. Tateda and K. Yamaguchi, *Int. J. Antimicrob. Agents*, 2007, **30**, 443–445.
- 27 B. Moya, L. Zamorano, C. Juan, Y. Ge and A. Oliver, *Antimicrob. Agents Chemother.*, 2010, **54**, 3933–3937.

- 28 N. Kohira, J. West, A. Ito, T. Ito-Horiyama, R. Nakamura, T. Sato, S. Rittenhouse, M. Tsuji and Y. Yamano, *Antimicrob. Agents Chemother.*, 2015, **60**, 729–734.
- 29 A. Ito, N. Kohira, S. K. Bouchillon, J. West, S. Rittenhouse, H. S. Sader, P. R. Rhomberg, R. N. Jones, H. Yoshizawa, R. Nakamura, M. Tsuji and Y. Yamano, *The Journal of antimicrobial chemotherapy*, 2016, **71**, 670–677.
- 30 U. Mollmann, L. Heinisch, A. Bauernfeind, T. Kohler and D. Ankel-Fuchs, *BioMetals*, 2009, **22**, 615–624.
- 31 Y. Ishii, K. Tateda, T. Horiyama, M. Tsuji, Y. Yamano, J. Shimada and K. Yamaguchi, presented in part at the 54th *Interscience Conference on Antimicrobial Agents and Chemotherapy*, Washington, DC, 2014.
- 32 ClinicalTrials.gov, Accessed 2/24/2016.
- 33 M. Babic, A. M. Hujer and R. A. Bonomo, *Drug Resist. Updates*, 2006, **9**, 142–156.
- 34 H. Knothe, P. Shah, V. Krcmery, M. Antal and S. Mitsuhashi, *Infection*, 1983, **11**, 315–317.
- 35 G. Cornaglia, H. Giamarellou and G. M. Rossolini, *Lancet Infect. Dis.*, 2011, **11**, 381–393.
- 36 A. Philippon, R. Labia and G. Jacoby, *Antimicrob. Agents Chemother.*, 1989, **33**, 1131–1136.
- 37 R. A. Powers, *Curr. Drug Targets*, 2015, **17**, 1051–1060.
- 38 D. M. Livermore, *Eur. J. Clin. Microbiol.*, 1987, **6**, 439–445.
- 39 N. Gupta, B. M. Limbago, J. B. Patel and A. J. Kallen, *Clin. Infect. Dis.*, 2011, **53**, 60–67.
- 40 H. Yigit, A. M. Queenan, G. J. Anderson, A. Domenech-Sanchez, J. W. Biddle, C. D. Steward, S. Alberti, K. Bush and F. C. Tenover, *Antimicrob. Agents Chemother.*, 2001, **45**, 1151–1161.
- 41 T. R. Walsh, *Clin. Microbiol. Infect.*, 2005, **11**(Suppl 6), 2–9.
- 42 P. Nordmann, L. Poirel, T. R. Walsh and D. M. Livermore, *Trends Microbiol.*, 2011, **19**, 588–595.
- 43 E. J. Zasowski, J. M. Rybak and M. J. Rybak, *Pharmacotherapy*, 2015, **35**, 755–770.
- 44 G. G. Zhanel, C. D. Lawson, H. Adam, F. Schweizer, S. Zelenitsky, P. R. Lagace-Wiens, A. Denisuk, E. Rubinstein, A. S. Gin, D. J. Hoban, J. P. Lynch 3rd and J. A. Karlowsky, *Drugs*, 2013, **73**, 159–177.
- 45 H. Li, M. Estabrook, G. A. Jacoby, W. W. Nichols, R. T. Testa and K. Bush, *Antimicrob. Agents Chemother.*, 2015, **59**, 1789–1793.
- 46 A. Yoshizumi, Y. Ishii, K. Aoki, R. Testa, W. W. Nichols and K. Tateda, *J. Infect. Chemother.*, 2015, **21**, 148–151.
- 47 H. S. Sader, M. Castanheira, R. E. Mendes, R. K. Flamm, D. J. Farrell and R. N. Jones, *Antimicrob. Agents Chemother.*, 2015, **59**, 3656–3659.
- 48 H. S. Sader, M. Castanheira, R. K. Flamm, D. J. Farrell and R. N. Jones, *Antimicrob. Agents Chemother.*, 2014, **58**, 1684–1692.
- 49 L. J. Dubreuil, S. Mahieux, C. Neut, C. Miossec and J. Pace, *Int. J. Antimicrob. Agents*, 2012, **39**, 500–504.
- 50 D. M. Citron, K. L. Tyrrell, V. Merriam and E. J. Goldstein, *Antimicrob. Agents Chemother.*, 2011, **55**, 3616–3620.
- 51 S. D. Lahiri, M. R. Johnstone, P. L. Ross, R. E. McLaughlin, N. B. Olivier and R. A. Alm, *Antimicrob. Agents Chemother.*, 2014, **58**, 5704–5713.
- 52 S. D. Lahiri, S. Mangani, T. Durand-Reville, M. Benvenuti, F. De Luca, G. Sanyal and J. D. Docquier, *Antimicrob. Agents Chemother.*, 2013, **57**, 2496–2505.
- 53 S. D. Lahiri, S. Mangani, H. Jahic, M. Benvenuti, T. F. Durand-Reville, F. De Luca, D. E. Ehmann, G. M. Rossolini, R. A. Alm and J. D. Docquier, *ACS Chem. Biol.*, 2015, **10**, 591–600.
- 54 N. P. Krishnan, N. Q. Nguyen, K. M. Papp-Wallace, R. A. Bonomo and F. van den Akker, *PLoS One*, 2015, **10**, e0136813.
- 55 A. Bonnefoy, C. Dupuis-Hamelin, V. Steier, C. Delachaume, C. Seys, T. Stachyra, M. Fairley, M. Guitton and M. Lampilas, *J. Antimicrob. Chemother.*, 2004, **54**, 410–417.
- 56 D. E. Ehmann, H. Jahic, P. L. Ross, R. F. Gu, J. Hu, G. Kern, G. K. Walkup and S. L. Fisher, *Proc. Natl. Acad. Sci. U. S. A.*, 2012, **109**, 11663–11668.
- 57 Z. Aktas, C. Kayacan and O. Oncul, *Int. J. Antimicrob. Agents*, 2012, **39**, 86–89.
- 58 K. Young, S. Raghoobar and N. Hairston, *et al.*, presented in part at the 50th *Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC)*, Boston, MA, 2010.
- 59 D. M. Livermore, M. Warner and S. Mushtaq, *J. Antimicrob. Chemother.*, 2013, **68**, 2286–2290.
- 60 E. B. Hirsch, K. R. Ledesma, K. T. Chang, M. S. Schwartz, M. R. Motyl and V. H. Tam, *Antimicrob. Agents Chemother.*, 2012, **56**, 3753–3757.
- 61 K. Young, M. Hackel and C. Lascols, *et al.*, presented in part at the 52nd *Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC)*, San Francisco, CA, 2012.
- 62 A. Lapuebla, M. Abdallah, O. Olafisoye, C. Cortes, C. Urban, D. Landman and J. Quale, *Antimicrob. Agents Chemother.*, 2015, **59**, 5029–5031.
- 63 T. A. Blizzard, H. Chen, S. Kim, J. Wu, R. Bodner, C. Gude, J. Imbriglio, K. Young, Y. W. Park, A. Ogawa, S. Raghoobar, N. Hairston, R. E. Painter, D. Wisniewski, G. Scapin, P. Fitzgerald, N. Sharma, J. Lu, S. Ha, J. Hermes and M. L. Hammond, *Bioorg. Med. Chem. Lett.*, 2014, **24**, 780–785.
- 64 E. Wenzler, M. H. Gotfried, J. S. Loutit, S. Durso, D. C. Griffith, M. N. Dudley and K. A. Rodvold, *Antimicrob. Agents Chemother.*, 2015, **59**, 7232–7239.
- 65 A. Lapuebla, M. Abdallah, O. Olafisoye, C. Cortes, C. Urban, J. Quale and D. Landman, *Antimicrob. Agents Chemother.*, 2015, **59**, 4856–4860.
- 66 D. M. Livermore and S. Mushtaq, *J. Antimicrob. Chemother.*, 2013, **68**, 1825–1831.
- 67 E. J. Goldstein, D. M. Citron, K. L. Tyrrell and C. V. Merriam, *Antimicrob. Agents Chemother.*, 2013, **57**, 2620–2630.
- 68 R. Smoum, A. Rubinstein, V. M. Dembitsky and M. Srebnik, *Chem. Rev.*, 2012, **112**, 4156–4220.
- 69 P. A. Kiener and S. G. Waley, *Biochem. J.*, 1978, **169**, 197–204.
- 70 O. Dobozy, I. Mile, I. Ferencz and V. Csanyi, *Acta Biochim. Biophys. Acad. Sci. Hung.*, 1971, **6**, 97–105.

- 71 N. C. Strynadka, H. Adachi, S. E. Jensen, K. Johns, A. Sielecki, C. Betzel, K. Sutoh and M. N. James, *Nature*, 1992, **359**, 700–705.
- 72 R. Martin and J. B. Jones, *Tetrahedron Lett.*, 1995, **36**, 8399–8402.
- 73 N. C. Strynadka, R. Martin, S. E. Jensen, M. Gold and J. B. Jones, *Nat. Struct. Biol.*, 1996, **3**, 688–695.
- 74 S. Ness, R. Martin, A. M. Kindler, M. Paetzel, M. Gold, S. E. Jensen, J. B. Jones and N. C. Strynadka, *Biochemistry*, 2000, **39**, 5312–5321.
- 75 F. Morandi, E. Caselli, S. Morandi, P. J. Focia, J. Blazquez, B. K. Shoichet and F. Prati, *J. Am. Chem. Soc.*, 2003, **125**, 685–695.
- 76 X. Wang, G. Minasov, J. Blazquez, E. Caselli, F. Prati and B. K. Shoichet, *Biochemistry*, 2003, **42**, 8434–8444.
- 77 S. J. Hecker, K. R. Reddy, M. Totrov, G. C. Hirst, O. Lomovskaya, D. C. Griffith, P. King, R. Tsivkovski, D. Sun, M. Sabet, Z. Tarazi, M. C. Clifton, K. Atkins, A. Raymond, K. T. Potts, J. Abendroth, S. H. Boyer, J. S. Loutit, E. E. Morgan, S. Durso and M. N. Dudley, *J. Med. Chem.*, 2015, **58**, 3682–3692.
- 78 R. C. Moellering, *Ann. Intern. Med.*, 2003, **138**, 135–142.
- 79 C. W. Ford, G. E. Zurenko and M. R. Barbachyn, *Curr. Drug Targets: Infect. Disord.*, 2001, **1**, 181–199.
- 80 A. M. Slee, M. A. Wuonola, R. J. McRipley, I. Zajac, M. J. Zawada, P. T. Bartholomew, W. A. Gregory and M. Forbes, *Antimicrob. Agents Chemother.*, 1987, **31**, 1791–1797.
- 81 M. R. Barbachyn and C. W. Ford, *Angew. Chem., Int. Ed.*, 2003, **42**, 2010–2023.
- 82 W. A. Gregory, D. R. Brittelli, C. L. Wang, H. S. Kezar, 3rd, R. K. Carlson, C. H. Park, P. F. Corless, S. J. Miller, P. Rajagopalan and M. A. Wuonola, *et al.*, *J. Med. Chem.*, 1990, **33**, 2569–2578.
- 83 W. A. Gregory, D. R. Brittelli, C. L. Wang, M. A. Wuonola, R. J. McRipley, D. C. Eustice, V. S. Eberly, P. T. Bartholomew, A. M. Slee and M. Forbes, *J. Med. Chem.*, 1989, **32**, 1673–1681.
- 84 C. H. Park, D. R. Brittelli, C. L. Wang, F. D. Marsh, W. A. Gregory, M. A. Wuonola, R. J. McRipley, V. S. Eberly, A. M. Slee and M. Forbes, *J. Med. Chem.*, 1992, **35**, 1156–1165.
- 85 M. R. Barbachyn, G. J. Cleek, L. A. Dolak, S. A. Garmon, J. Morris, E. P. Seest, R. C. Thomas, D. S. Toops, W. Watt, D. G. Wishka, C. W. Ford, G. E. Zurenko, J. C. Hamel, R. D. Schaadt, D. Stapert, B. H. Yagi, W. J. Adams, J. M. Friis, J. G. Slatte, J. P. Sams, N. L. Oien, M. J. Zaya, L. C. Wienkers and M. A. Wynalda, *J. Med. Chem.*, 2003, **46**, 284–302.
- 86 A. D. Borthwick, K. Biggedike, V. Rocherolle, D. M. Cox and G. A. C. Chung, *Med. Chem. Res.*, 1996, **6**, 22–27.
- 87 A. Dennis and T. Villette, *Bioorg. Med. Chem. Lett.*, 1994, **4**, 1925–1930.
- 88 C. A. Quesnelle, P. Gill, S. Roy, M. Dodier, A. Marinier, A. Martel, L. B. Snyder, S. V. D'Andrea, J. J. Bronson, M. Frosco, D. Beaulieu, G. A. Warr, K. L. Denbleyker, T. M. Stickle, H. Yang, S. E. Chaniewski, C. A. Ferraro, D. Taylor, J. W. Russell, K. S. Santone, J. Clarke, R. L. Drain, J. O. Knipe, K. Mosure and J. F. Barrett, *Bioorg. Med. Chem. Lett.*, 2005, **15**, 2728–2733.
- 89 L. B. Snyder, Z. Meng, R. Mate, S. V. D'Andrea, A. Marinier, C. A. Quesnelle, P. Gill, K. L. Denbleyker, J. C. Fung-Tomc, M. Frosco, A. Martel, J. F. Barrett and J. J. Bronson, *Bioorg. Med. Chem. Lett.*, 2004, **14**, 4735–4739.
- 90 S. J. Brickner, *Curr. Pharm. Des.*, 1996, **2**, 175–194.
- 91 M. R. Barbachyn, D. K. Hutchinson, S. J. Brickner, M. H. Cynamon, J. O. Kilburn, S. P. Klemens, S. E. Glickman, K. C. Grega, S. K. Hendges, D. S. Toops, C. W. Ford and G. E. Zurenko, *J. Med. Chem.*, 1996, **39**, 680–685.
- 92 Y. W. Jo, W. B. Im, J. K. Rhee, M. J. Shim, W. B. Kim and E. C. Choi, *Bioorg. Med. Chem.*, 2004, **12**, 5909–5915.
- 93 M. B. Gravestock, D. G. Acton, M. J. Betts, M. Dennis, G. Hatter, A. McGregor, M. L. Swain, R. G. Wilson, L. Woods and A. Wookey, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 4179–4186.
- 94 O. A. Phillips, E. E. Udo, A. A. Ali and N. Al-Hassawi, *Bioorg. Med. Chem.*, 2003, **11**, 35–41.
- 95 S. M. Swaney, H. Aoki, M. C. Ganoza and D. L. Shinabarger, *Antimicrob. Agents Chemother.*, 1998, **42**, 3251–3255.
- 96 J. J. Kisgen, H. Mansour, N. R. Unger and L. M. Childs, *Am. J. Health-Syst. Pharm.*, 2014, **71**, 621–633.
- 97 J. B. Locke, G. E. Zurenko, K. J. Shaw and K. Bartizal, *Clin. Infect. Dis.*, 2014, **58**(Suppl 1), S35–42.
- 98 G. E. Zurenko, J. K. Gibson, D. L. Shinabarger, P. A. Aristoff, C. W. Ford and W. G. Tarpley, *Curr. Opin. Pharmacol.*, 2001, **1**, 470–476.
- 99 A. Schumacher, R. Trittler, J. A. Bohnert, K. Kummerer, J. M. Pages and W. V. Kern, *J. Antimicrob. Chemother.*, 2007, **59**, 1261–1264.
- 100 M. C. Roberts, *Curr. Drug Targets: Infect. Disord.*, 2004, **4**, 207–215.
- 101 M. C. Roberts, *FEMS Microbiol. Lett.*, 2008, **282**, 147–159.
- 102 G. G. Zhanel, R. Love, H. Adam, A. Golden, S. Zelenitsky, F. Schweizer, B. Gorityala, P. R. Lagace-Wiens, E. Rubinstein, A. Walkty, A. S. Gin, M. Gilmour, D. J. Hoban, J. P. Lynch, 3rd and J. A. Karlowsky, *Drugs*, 2015, **75**, 253–270.
- 103 M. E. Stryjewski and G. R. Corey, *Clin. Infect. Dis.*, 2014, **58**(Suppl 1), S10–19.
- 104 J. B. Locke, D. E. Zuill, C. R. Scharn, J. Deane, D. F. Sahm, G. A. Denys, R. V. Goering and K. J. Shaw, *Antimicrob. Agents Chemother.*, 2014, **58**, 6592–6598.
- 105 G. Morales, J. J. Picazo, E. Baos, F. J. Candel, A. Arribi, B. Pelaez, R. Andrade, M. A. de la Torre, J. Fereres and M. Sanchez-Garcia, *Clin. Infect. Dis.*, 2010, **50**, 821–825.
- 106 E. B. Chahine, A. J. Sucher and S. D. Knutsen, *Consult. Pharm.*, 2015, **30**, 386–394.
- 107 M. J. Durkin and G. R. Corey, *Ther. Clin. Risk Manage.*, 2015, **11**, 857–862.
- 108 R. G. Hall 2nd and H. N. Michaels, *Infect. Drug Resist.*, 2015, **8**, 75–82.
- 109 L. Morata, J. Mensa and A. Soriano, *Curr. Opin. Pharmacol.*, 2015, **24**, 45–51.
- 110 Z. A. Kanafani and G. R. Corey, *Expert Opin. Invest. Drugs*, 2012, **21**, 515–522.

- 111 K. Shen, K. Yoshikawa, T. Tanaka and *et al.*, presented in part at the 43rd *Society of Critical Care Medicine Critical Care Congress*, San Francisco, CA, 2014.
- 112 T. Tanaka, Y. Hayashi, K. Okumura and *et al.*, presented in part at the 24th *European Congress of Clinical Microbiology and Infectious Diseases*, Barcelona, Spain, 2014.
- 113 J. B. Locke, J. Finn, M. Hilgers, G. Morales, S. Rahawi, G C. K., J. J. Picazo, W. Im, K. J. Shaw and J. L. Stein, *Antimicrob. Agents Chemother.*, 2010, 54, 5337–5343.
- 114 J. Locke, J. Finn, M. Hilgers and *et al.*, presented in part at the 50th *Interscience Conference on Antimicrobial Agents and Chemotherapy*, Boston, MA, 2010.
- 115 K. Michalska, I. Karpiuk, M. Krol and S. Tyski, *Bioorg. Med. Chem.*, 2013, 21, 577–591.
- 116 K. J. Shaw, S. Poppe, R. Schaadt, V. Brown-Driver, J. Finn, C. M. Pillar, D. Shinabarger and G. Zurenko, *Antimicrob. Agents Chemother.*, 2008, 52, 4442–4447.
- 117 J. H. Yum, S. H. Choi, D. Yong, Y. Chong, W. B. Im, D. K. Rhee and K. Lee, *Antimicrob. Agents Chemother.*, 2010, 54, 5381–5386.
- 118 P. Prokocimer, P. Bien, C. Deanda, C. M. Pillar and K. Bartizal, *Antimicrob. Agents Chemother.*, 2012, 56, 4608–4613.
- 119 S. D. Brown and M. M. Traczewski, *Antimicrob. Agents Chemother.*, 2010, 54, 2063–2069.
- 120 R. Schaadt, D. Sweeney, D. Shinabarger and G. Zurenko, *Antimicrob. Agents Chemother.*, 2009, 53, 3236–3239.
- 121 H. Chen, Q. Yang, R. Zhang, W. He, X. Ma, J. Zhang, F. Xia, F. Zhao, J. Cao, Y. Liu, W. Wu, D. Hu, Q. Wang, C. Zhao, F. Zhang, X. Wang, Z. Wang, H. Li and H. Wang, *Int. J. Antimicrob. Agents*, 2014, 44, 276–277.
- 122 R. N. Jones, G. J. Moet, H. S. Sader, R. E. Mendes and M. Castanheira, *J. Antimicrob. Chemother.*, 2009, 63, 716–720.
- 123 K. J. Shaw and M. R. Barbachyn, *Ann. N. Y. Acad. Sci.*, 2011, 1241, 48–70.
- 124 J. Ippolito, B. Kanyo, B. Wimberly and *et al.*, presented in part at the 45th *Interscience Conference on Antimicrobial Agents and Chemotherapy*, Washington, DC, 2005.
- 125 D. Wang, E. Sherer and E. Duffy, presented in part at the 45th *Interscience Conference on Antimicrobial Agents and Chemotherapy*, Washington, DC, 2005.
- 126 J. Zhou, A. Bhattacharjee, S. Chen, Y. Chen, E. Duffy, J. Farmer, J. Goldberg, R. Hanselmann, J. A. Ippolito, R. Lou, A. Orbin, A. Oyelere, J. Salvino, D. Springer, J. Tran, D. Wang, Y. Wu and G. Johnson, *Bioorg. Med. Chem. Lett.*, 2008, 18, 6179–6183.
- 127 J. Zhou, A. Bhattacharjee, S. Chen, Y. Chen, E. Duffy, J. Farmer, J. Goldberg, R. Hanselmann, J. A. Ippolito, R. Lou, A. Orbin, A. Oyelere, J. Salvino, D. Springer, J. Tran, D. Wang, Y. Wu and G. Johnson, *Bioorg. Med. Chem. Lett.*, 2008, 18, 6175–6178.
- 128 L. Lawrence, P. Danese, J. DeVito, F. Franceschi and J. Sutcliffe, *Antimicrob. Agents Chemother.*, 2008, 52, 1653–1662.
- 129 E. Skripkin, T. S. McConnell, J. DeVito, L. Lawrence, J. A. Ippolito, E. M. Duffy, J. Sutcliffe and F. Franceschi, *Antimicrob. Agents Chemother.*, 2008, 52, 3550–3557.
- 130 M. F. Gordeev and Z. Y. Yuan, *J. Med. Chem.*, 2014, 57, 4487–4497.
- 131 C. R. Li, Q. Q. Zhai, X. K. Wang, X. X. Hu, G. Q. Li, W. X. Zhang, J. Pang, X. Lu, H. Yuan, M. F. Gordeev, L. T. Chen, X. Y. Yang and X. F. You, *Antimicrob. Agents Chemother.*, 2014, 58, 2418–2421.
- 132 A. Kali, M. V. Charles and S. Srirangaraj, *Australas Med J*, 2015, 8, 253–262.
- 133 M. F. Gordeev, C. Hackbarth, M. R. Barbachyn, L. S. Banitt, J. R. Gage, G. W. Luehr, M. Gomez, J. Trias, S. E. Morin, G. E. Zurenko, C. N. Parker, J. M. Evans, R. J. White and D. V. Patel, *Bioorg. Med. Chem. Lett.*, 2003, 13, 4213–4216.
- 134 C. Hubschwerlen, J. L. Specklin, D. K. Baeschlin, Y. Borer, S. Haefeli, C. Sigwalt, S. Schroeder and H. H. Locher, *Bioorg. Med. Chem. Lett.*, 2003, 13, 4229–4233.
- 135 C. Hubschwerlen, J. L. Specklin, C. Sigwalt, S. Schroeder and H. H. Locher, *Bioorg. Med. Chem.*, 2003, 11, 2313–2319.
- 136 H. H. Locher, P. Caspers, T. Bruyere, S. Schroeder, P. Pfaff, A. Knezevic, W. Keck and D. Ritz, *Antimicrob. Agents Chemother.*, 2014, 58, 901–908.
- 137 M. U. Rashid, A. Dalhoff, A. Weintraub and C. E. Nord, *Anaerobe*, 2014, 28, 216–219.
- 138 M. U. Rashid, H. M. Lozano, A. Weintraub and C. E. Nord, *Anaerobe*, 2013, 20, 32–35.
- 139 H. H. Locher, P. Seiler, X. Chen, S. Schroeder, P. Pfaff, M. Enderlin, A. Klenk, E. Fournier, C. Hubschwerlen, D. Ritz, C. P. Kelly and W. Keck, *Antimicrob. Agents Chemother.*, 2014, 58, 892–900.
- 140 D. Baldoni, M. Gutierrez, W. Timmer and J. Dingemanse, *J. Antimicrob. Chemother.*, 2014, 69, 706–714.
- 141 A. A. Miller, G. L. Bundy, J. E. Mott, J. E. Skepner, T. P. Boyle, D. W. Harris, A. E. Hromockyj, K. R. Marotti, G. E. Zurenko, J. B. Munzner, M. T. Sweeney, G. F. Bammert, J. C. Hamel, C. W. Ford, W. Z. Zhong, D. R. Graber, G. E. Martin, F. Han, L. A. Dolak, E. P. Seest, J. C. Ruble, G. M. Kamilar, J. R. Palmer, L. S. Banitt, A. R. Hurd and M. R. Barbachyn, *Antimicrob. Agents Chemother.*, 2008, 52, 2806–2812.
- 142 G. S. Basarab, G. H. Kern, J. McNulty, J. P. Mueller, K. Lawrence, K. Vishwanathan, R. A. Alm, K. Barvian, P. Doig, V. Galullo, H. Gardner, M. Gowravaram, M. Huband, A. Kimzey, M. Morningstar, A. Kutschke, S. D. Lahiri, M. Perros, R. Singh, V. J. Schuck, R. Tommasi, G. Walkup and J. V. Newman, *Sci. Rep.*, 2015, 5, 11827.
- 143 M. Unemo, J. Ringlander, C. Wiggins, H. Fredlund, S. Jacobsson and M. Cole, *Antimicrob. Agents Chemother.*, 2015, 59, 5220–5225.
- 144 M. D. Huband, P. A. Bradford, L. G. Otterson, G. S. Basarab, A. C. Kutschke, R. A. Giacobbe, S. A. Patey, R. A. Alm, M. R. Johnstone, M. E. Potter, P. F. Miller and J. P. Mueller, *Antimicrob. Agents Chemother.*, 2015, 59, 467–474.
- 145 D. J. Biedenbach, S. K. Bouchillon, M. Hackel, L. A. Miller, N. E. Scangarella-Oman, C. Jakielaszek and D. F. Sahm, *Antimicrob. Agents Chemother.*, 2016, 60, 1918–1923.
- 146 S. B. Singh, *Bioorg. Med. Chem. Lett.*, 2014, 24, 3683–3689.

- 147 B. D. Bax, P. F. Chan, D. S. Eggleston, A. Fosberry, D. R. Gentry, F. Gorrec, I. Giordano, M. M. Hann, A. Hennessy, M. Hibbs, J. Huang, E. Jones, J. Jones, K. K. Brown, C. J. Lewis, E. W. May, M. R. Saunders, O. Singh, C. E. Spitzfaden, C. Shen, A. Shillings, A. J. Theobald, A. Wohlkonig, N. D. Pearson and M. N. Gwynn, *Nature*, 2010, **466**, 935–940.
- 148 K. Widdowson and A. Hennessy, *Future Med. Chem.*, 2010, **2**, 1619–1622.
- 149 S. Lemaire, F. Van Bambeke and P. M. Tulkens, *Int. J. Antimicrob. Agents*, 2011, **38**, 52–59.
- 150 S.-E. Wohlert, T. Jaetsch, B. Gallenkamp, H. J. Knops, N. Lui, M. Preiss, D. Haebich and H. Labischinski, presented in part at the 48th *Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC)*, Washington, D.C., 2008.
- 151 H. J. Adam, N. M. Laing, C. R. King, B. Lulashnyk, D. J. Hoban and G. G. Zhanel, *Antimicrob. Agents Chemother.*, 2009, **53**, 4915–4920.
- 152 X. Qin and H. Huang, *Drug Des., Dev. Ther.*, 2014, **8**, 765–774.
- 153 Z. X. Li, Y. N. Liu, R. Wang and A. M. Li, *Clin. Ter.*, 2015, **166**, e374–380.
- 154 Y. H. Chen, C. Y. Liu, J. J. Lu, C. H. King and P. R. Hsueh, *J. Antimicrob. Chemother.*, 2009, **64**, 1226–1229.
- 155 A. M. Nilius, L. L. Shen, D. Hensey-Rudloff, L. S. Almer, J. M. Beyer, D. J. Balli, Y. Cai and R. K. Flamm, *Antimicrob. Agents Chemother.*, 2003, **47**, 3260–3269.
- 156 F. Van Bambeke, *Future Microbiol.*, 2015, **10**, 1111–1123.
- 157 S. Lemaire, P. M. Tulkens and F. Van Bambeke, *Antimicrob. Agents Chemother.*, 2011, **55**, 649–658.
- 158 W. So, J. L. Crandon and D. P. Nicolau, *J. Urol.*, 2015, **194**, 563–570.
- 159 E. M. Duffy, J. A. DeVito, J. Remy and E. S. Burak, presented in part at the 50th *Interscience Conference on Antimicrobial Agents and Chemotherapy (ICAAC)*, Boston, MA, 2010.
- 160 D. E. Brodersen, W. M. Clemons Jr., A. P. Carter, R. J. Morgan-Warren, B. T. Wimberly and V. Ramakrishnan, *Cell*, 2000, **103**, 1143–1154.
- 161 P. E. Sum and P. Petersen, *Bioorg. Med. Chem. Lett.*, 1999, **9**, 1459–1462.
- 162 J. Bergeron, M. Ammirati, D. Danley, L. James, M. Norcia, J. Retsema, C. A. Strick, W. G. Su, J. Sutcliffe and L. Wondrack, *Antimicrob. Agents Chemother.*, 1996, **40**, 2226–2228.
- 163 P. J. Petersen, N. V. Jacobus, W. J. Weiss, P. E. Sum and R. T. Testa, *Antimicrob. Agents Chemother.*, 1999, **43**, 738–744.
- 164 M. W. Garrison, J. J. Neumiller and S. M. Setter, *Clin. Ther.*, 2005, **27**, 12–22.
- 165 T. C. Barden, B. L. Buckwalter, R. T. Testa, P. J. Petersen and V. J. Lee, *J. Med. Chem.*, 1994, **37**, 3205–3211.
- 166 G. M. Eliopoulos, C. B. Wennersten, G. Cole and R. C. Moellering, *Antimicrob. Agents Chemother.*, 1994, **38**, 534–541.
- 167 L. Jenner, A. L. Starosta, D. S. Terry, A. Mikolajka, L. Filonava, M. Yusupov, S. C. Blanchard, D. N. Wilson and G. Yusupova, *Proc. Natl. Acad. Sci. U. S. A.*, 2013, **110**, 3812–3816.
- 168 A. B. Maccone, B. K. Caruso, R. G. Leahy, J. Donatelli, S. Weir, M. P. Draper, S. K. Tanaka and S. B. Levy, *Antimicrob. Agents Chemother.*, 2014, **58**, 1127–1135.
- 169 L. Honeyman, M. Ismail, M. L. Nelson, B. Bhatia, T. E. Bowser, J. Chen, R. Mechiche, K. Ohemeng, A. K. Verma, E. P. Cannon, A. Maccone, S. K. Tanaka and S. Levy, *Antimicrob. Agents Chemother.*, 2015, **59**, 7044–7053.
- 170 X. Y. Xiao, D. K. Hunt, J. Zhou, R. B. Clark, N. Dunwoody, C. Fyfe, T. H. Grossman, W. J. O'Brien, L. Plamondon, M. Ronn, C. Sun, W. Y. Zhang and J. A. Sutcliffe, *J. Med. Chem.*, 2012, **55**, 597–605.
- 171 A. Cammarata and S. J. Yau, *J. Med. Chem.*, 1970, **13**, 93–97.
- 172 J. A. Sutcliffe, W. O'Brien, C. Fyfe and T. H. Grossman, *Antimicrob. Agents Chemother.*, 2013, **57**, 5548–5558.
- 173 T. H. Grossman, A. L. Starosta, C. Fyfe, W. O'Brien, D. M. Rothstein, A. Mikolajka, D. N. Wilson and J. A. Sutcliffe, *Antimicrob. Agents Chemother.*, 2012, **56**, 2559–2564.
- 174 G. G. Zhanel, D. Cheung, H. Adam, S. Zelenitsky, A. Golden, F. Schweizer, B. Gorityala, P. R. Lagace-Wiens, A. Walkty, A. S. Gin, D. J. Hoban and J. A. Karlowsky, *Drugs*, 2016, **76**, 567–588.
- 175 S. Sivapalasingam and N. H. Steigbigel, in *Mandell, Douglas, and Bennett's Principles and Practice of Infectious Diseases*, ed. J. E. Bennett, R. Dolin and M. J. Blaser, Saunders - Elsevier, Inc., Philadelphia, PA, 8th edn, 2014, ch. 29, vol. 1, pp. 358–376.
- 176 A. Bryskier, *Clin. Microbiol. Infect.*, 2000, **6**, 661–669.
- 177 H. Okamoto, S. Miyazaki, K. Tateda, Y. Ishii and K. Yamaguchi, *J. Antimicrob. Chemother.*, 2000, **46**, 797–802.
- 178 R. Chollet, J. Chevalier, A. Bryskier and J. M. Pages, *Antimicrob. Agents Chemother.*, 2004, **48**, 3621–3624.
- 179 L. Xiong, S. Shah, P. Mauvais and A. S. Mankin, *Mol. Microbiol.*, 1999, **31**, 633–639.
- 180 F. Blasi, D. J. Farrell and L. Dubreuil, *Diagn. Microbiol. Infect. Dis.*, 2009, **63**, 302–308.
- 181 D. Felmingham and D. J. Farrell, *J. Infect.*, 2006, **52**, 178–180.
- 182 M. R. Hammerschlag, P. M. Roblin and C. M. Bebear, *J. Antimicrob. Chemother.*, 2001, **48**(Suppl T1), 25–31.
- 183 N. H. Georgopapadakou, *Expert Opin. Invest. Drugs*, 2014, **23**, 1313–1319.
- 184 P. S. Barie, *Surg. Infect.: Sel. Antibiot. Ther.*, 2006, **7**, 247–249.
- 185 A. W. Mathews, *Wall St. J., East. Ed.*, 2006, A1–A12.
- 186 H. Mansour, E. B. Chahine, L. R. Karaoui and R. M. El-Lababidi, *Ann. Pharmacother.*, 2013, **47**, 368–379.
- 187 E. C. Kouvela, D. L. Kalpaxis, D. N. Wilson and G. P. Dinios, *Antimicrob. Agents Chemother.*, 2009, **53**, 1411–1419.
- 188 A. M. Nilius and Z. Ma, *Curr. Opin. Pharmacol.*, 2002, **2**, 493–500.
- 189 C. M. Barrera, A. Mykietiuks, H. Metev, M. F. Nitu, N. Karimjee, P. A. Doreski, I. Mitha, C. M. Tanaseanu, J. M. Molina, Y. Antonovsky, D. J. Van Rensburg, B. H. Rowe, J. Flores-Figueroa, B. Rewerska, K. Clark, K. Keedy, A. Sheets,

- D. Scott, G. Horwith, A. F. Das, B. Jamieson, P. Fernandes, D. Oldach and S.-O. P. Team, *Lancet Infect. Dis.*, 2016, **16**, 421–430.
- 190 F. Van Bambeke and P. M. Tulkens, *Expert Rev. Anti-Infect. Ther.*, 2016, **14**, 311–324.
- 191 P. McGhee, C. Clark, K. M. Kosowska-Shick, K. Nagai, B. Dewasse, L. Beachel and P. C. Appelbaum, *Antimicrob. Agents Chemother.*, 2010, **54**, 230–238.
- 192 B. Llano-Sotelo, J. Dunkle, D. Klepacki, W. Zhang, P. Fernandes, J. H. Cate and A. S. Mankin, *Antimicrob. Agents Chemother.*, 2010, **54**, 4961–4970.
- 193 G. G. Zhanel, C. D. Lawson, S. Zelenitsky, B. Findlay, F. Schweizer, H. Adam, A. Walkty, E. Rubinstein, A. S. Gin, D. J. Hoban, J. P. Lynch and J. A. Karlowksy, *Expert Rev. Anti-Infect. Ther.*, 2012, **10**, 459–473.
- 194 J. B. Aggen, E. S. Armstrong, A. A. Goldblum, P. Dozzo, M. S. Linsell, M. J. Gliedt, D. J. Hildebrandt, L. A. Feeney, A. Kubo, R. D. Matias, S. Lopez, M. Gomez, K. B. Wlasichuk, R. Diokno, G. H. Miller and H. E. Moser, *Antimicrob. Agents Chemother.*, 2010, **54**, 4636–4642.
- 195 P. Schneider, S. Hawser and K. Islam, *Bioorg. Med. Chem. Lett.*, 2003, **13**, 4217–4221.
- 196 G. E. Dale, C. Broger, A. D'Arcy, P. G. Hartman, R. DeHoogt, S. Jolidon, I. Kompis, A. M. Labhardt, H. Langen, H. Locher, M. G. Page, D. Stuber, R. L. Then, B. Wipf and C. Oefner, *J. Mol. Biol.*, 1997, **266**, 23–30.
- 197 C. Oefner, M. Bandera, A. Haldimann, H. Laue, H. Schulz, S. Mukhija, S. Parisi, L. Weiss, S. Lociuero and G. E. Dale, *J. Antimicrob. Chemother.*, 2009, **63**, 687–698.
- 198 C. A. Sincak and J. M. Schmidt, *Ann. Pharmacother.*, 2009, **43**, 1107–1114.
- 199 H. Laue, T. Valensise, A. Seguin, S. Lociuero, K. Islam and S. Hawser, *Antimicrob. Agents Chemother.*, 2009, **53**, 4542–4544.
- 200 H. Laue, L. Weiss, A. Bernardi, S. Hawser, S. Lociuero and K. Islam, *J. Antimicrob. Chemother.*, 2007, **60**, 1391–1394.
- 201 S. W. White, J. Zheng, Y. M. Zhang and C. O. Rock, *Annu. Rev. Biochem.*, 2005, **74**, 791–831.
- 202 Y. M. Zhang, Y. J. Lu and C. O. Rock, *Lipids*, 2004, **39**, 1055–1060.
- 203 H. Marrakchi, Y. M. Zhang and C. O. Rock, *Biochem. Soc. Trans.*, 2002, **30**, 1050–1055.
- 204 R. J. Heath, J. R. Rubin, D. R. Holland, E. Zhang, M. E. Snow and C. O. Rock, *J. Biolumin. Chemilumin.*, 1999, **274**, 11110–11114.
- 205 S. Brinster, G. Lamberet, B. Staels, P. Trieu-Cuot, A. Gruss and C. Poyart, *Nature*, 2009, **458**, 83–86.
- 206 W. Balemans, N. Lounis, R. Gilissen, J. Guillemont, K. Simmen, K. Andries and A. Koul, *Nature*, 2010, **463**, E3–E4.
- 207 S. Brinster, G. Lamberet, B. Staels, P. Trieu-Cuot, A. Gruss and C. Poyart, *Nature*, 2010, **463**, E4–E5.
- 208 J. B. Parsons, M. W. Frank, C. Subramanian, P. Saenkham and C. O. Rock, *Proc. Natl. Acad. Sci. U. S. A.*, 2011, **108**, 15378–15383.
- 209 J. B. Parsons and C. O. Rock, *Curr. Opin. Microbiol.*, 2011, **14**, 544–549.
- 210 S. Mehboob, J. Song, K. E. Hevener, P. C. Su, T. Boci, L. Brubaker, L. Truong, T. Mistry, J. Deng, J. L. Cook, B. D. Santarsiero, A. K. Ghosh and M. E. Johnson, *Bioorg. Med. Chem. Lett.*, 2015, **25**, 1292–1296.
- 211 C. J. Zheng, M. J. Sohn, S. Lee and W. G. Kim, *PLoS One*, 2013, **8**, e78922.
- 212 K. E. Hevener, S. Mehboob, P. C. Su, K. Truong, T. Boci, J. Deng, M. Ghassemi, J. L. Cook and M. E. Johnson, *J. Med. Chem.*, 2012, **55**, 268–279.
- 213 S. Mehboob, K. E. Hevener, K. Truong, T. Boci, B. D. Santarsiero and M. E. Johnson, *J. Med. Chem.*, 2012, **55**, 5933–5941.
- 214 J. Schiebel, A. Chang, H. Lu, M. V. Baxter, P. J. Tonge and C. Kisker, *Structure*, 2012, **20**, 802–813.
- 215 V. Gerusz, A. Denis, F. Faivre, Y. Bonvin, M. Oxoby, S. Briet, G. LeFralliec, C. Oliveira, N. Desroy, C. Raymond, L. Peltier, F. Moreau, S. Escaich, V. Vongsouthi, S. Floquet, E. Drocourt, A. Walton, L. Prouvensier, M. Saccomani, L. Durant, J. M. Genevard, V. Sam-Sambo and C. Soulama-Mouze, *J. Med. Chem.*, 2012, **55**, 9914–9928.
- 216 N. Liu, J. E. Cummings, K. England, R. A. Slayden and P. J. Tonge, *J. Antimicrob. Chemother.*, 2011, **66**, 564–573.
- 217 P. B. Sampson, C. Picard, S. Handerson, T. E. McGrath, M. Domagala, A. Leeson, V. Romanov, D. E. Awrey, D. Thambipillai, E. Bardouniotis, N. Kaplan, J. M. Berman and H. W. Pauls, *Bioorg. Med. Chem. Lett.*, 2009, **19**, 5355–5358.
- 218 J. Ramnauth, M. D. Surman, P. B. Sampson, B. Forrest, J. Wilson, E. Freeman, D. D. Manning, F. Martin, A. Toro, M. Domagala, D. E. Awrey, E. Bardouniotis, N. Kaplan, J. Berman and H. W. Pauls, *Bioorg. Med. Chem. Lett.*, 2009, **19**, 5359–5362.
- 219 J. A. Karlowksy, N. Kaplan, B. Hafkin, D. J. Hoban and G. G. Zhanel, *Antimicrob. Agents Chemother.*, 2009, **53**, 3544–3548.
- 220 C. J. Zheng, M. J. Sohn and W. G. Kim, *J. Antimicrob. Chemother.*, 2009, **63**, 949–953.
- 221 H. Lu and P. J. Tonge, *Acc. Chem. Res.*, 2008, **41**, 11–20.
- 222 G. S. Dhillon, S. Kaur, R. Pulicharla, S. K. Brar, M. Cledon, M. Verma and R. Y. Surampalli, *Int. J. Environ. Res. Public Health*, 2015, **12**, 5657–5684.
- 223 L. M. McMurtry, M. Oethinger and S. B. Levy, *Nature*, 1998, **394**, 531–532.
- 224 R. K. Flamm, P. R. Rhomberg, N. Kaplan, R. N. Jones and D. J. Farrell, *Antimicrob. Agents Chemother.*, 2015, **59**, 2583–2587.
- 225 B. Hafkin, N. Kaplan and B. Murphy, *Antimicrob. Agents Chemother.*, 2015, **60**, 1695–1701.
- 226 F. Wittke, V. Nicolas-Metral, A. Menetrey and *et al.*, presented in part at the 55th *Interscience Conference on Antimicrobial Agents and Chemotherapy*, San Diego, CA, 2015.
- 227 N. Kaplan, M. Albert, D. Awrey, E. Bardouniotis, J. Berman, T. Clarke, M. Dorsey, B. Hafkin, J. Ramnauth, V. Romanov, M. B. Schmid, R. Thalakada, J. Yethon and H. W. Pauls, *Antimicrob. Agents Chemother.*, 2012, **56**, 5865–5874.
- 228 J. A. Karlowksy, N. M. Laing, T. Baudry, N. Kaplan, D. Vaughan, D. J. Hoban and G. G. Zhanel, *Antimicrob. Agents Chemother.*, 2007, **51**, 1580–1581.

- 229 T. Bogdanovich, C. Clark, K. Kosowska-Shick, B. Dewasse, P. McGhee and P. C. Appelbaum, *Antimicrob. Agents Chemother.*, 2007, **51**, 4191–4195.
- 230 H. S. Park, Y. M. Yoon, S. J. Jung, C. M. Kim, J. M. Kim and J. H. Kwak, *J. Antimicrob. Chemother.*, 2007, **60**, 568–574.
- 231 J. H. Yum, C. K. Kim, D. Yong, K. Lee, Y. Chong, C. M. Kim, J. M. Kim, S. Ro and J. M. Cho, *Antimicrob. Agents Chemother.*, 2007, **51**, 2591–2593.
- 232 J. Schiebel, A. Chang, S. Shah, Y. Lu, L. Liu, P. Pan, M. W. Hirschbeck, M. Tareilus, S. Eltschkner, W. Yu, J. E. Cummings, S. E. Knudson, G. R. Bommineni, S. G. Walker, R. A. Slayden, C. A. Sottriffer, P. J. Tonge and C. Kisker, *J. Biolumin. Chemilumin.*, 2014, **289**, 15987–16005.
- 233 allphasepharma.com, <http://allphasepharma.com/dir/2015/12/23/2205/qidp-antibiotics-2015-year-end-update/>, Accessed 2/10/2016.